#### PATENT APPLICATION

#### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Docket No: Q75926

Ryuichi MORISHITA, et al.

Appln. No.: 10/615,262

Group Art Unit: 1633

Confirmation No.: 5695

Examiner: Robert M. KELLY

Filed: July 9, 2003

For:

MEDICAMENT COMPRISING HGF GENE

### **DECLARATION UNDER 37 C.F.R. § 1.132**

Mail Stop Amendment Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450

Sir:

I, Ryuichi MORISHITA, hereby declare and state:

THAT I am a citizen of Japan, and reside at c/o 1-41-4, Senriyama-nishi, Suita-shi, Osaka, 565-0851, Japan;

THAT I am the first inventor of the above-identified patent application;

THAT I attach hereto an up to date copy of my curriculum vitae;

THAT I obtained the qualification of MD from Osaka University Medical School in Japan in 1987, and the qualification of Ph.D. from Osaka University Medical School in Japan in 1991.

THAT I currently hold a position as a professor of Division of Clinical Gene Therapy, Osaka University Medical School,

I further declare and state as follows:

I am familiar with the Office Action mailed June 13, 2007, in which the Examiner rejects Claims 7-11 on the ground of nonstatutory double patenting as claiming an invention that is not patentably distinct from the invention claimed in claims 1-3 of USP 6,936,594 (Morishita et al.).

The Examiner appears to assert that Morishita et al. teaches treating the cerebrum at the objective site, which includes the subarachnoid space, citing page 10, last paragraph. The Examiner seems to assert that if the brain were experiencing insufficiency of peripheral circulation, the subarachnoid space would be considered an area affected by the insufficiency, and thus would be the site of administration of the gene. Thus, the Examiner asserts that treating cerebrovascular disorders by administering the HGF gene to the subarachnoid space teaches or suggests treating insufficiency of peripheral circulation by administering the HGF gene to the affected area.

For the following reasons, I respectfully disagree with the Examiner.

Claims 1, 2, and 3 of Morishita et al. relate to "treatment of cardiovascular disorders," "treatment of reduced blood flow," and "promoting cerebral angiogenesis," respectively, by administering HGF gene to the subarachnoid space. However, "treatment of cardiovascular disorders," "treatment of reduced blood flow," and "promoting cerebral angiogenesis," do not teach or suggest treatment of insufficiency of peripheral circulation or peripheral angiostenosis, by injection of the HGF gene into the subarachnoid space.

More specifically, the claims of Morishita et al. relate to the treatment of blood disorders in the brain, namely, cerebrovascular disorders. In contrast, the presently claimed invention relates to the treatment of insufficiency of peripheral circulation or peripheral angiostenosis.

Attorney Docket No.: Q75926

U.S. Application No.: 10/615,262

One of ordinary skill in the art of medicine knows that brain is classified as part of the central nervous system. However, the term "central" is the antonym of the term "peripheral." My opinion is supported by Steadman's Medical Dictionary, 28th edition, a copy of which is submitted herewith. Steadman's Medical Dictionary defines the "brain" as "That part of the central nervous system contained within the cranium" (page 250, right column). In contrast, Steadman's Medical Dictionary defines "peripheral" as "opposite of central (centralis)" (page 1463, right column).

Therefore, in my opinion, the cerebrovascular disorders of Morishita et al. have no relevance to the insufficiency of peripheral circulation or peripheral angiostenosis of the present claims, and, as a result, the claims of Morishita et al. do not teach or suggest the presently claimed invention.

I declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date: 2007/ Dec 05

Ryuichi MORISHITA

## CURRICULUM VITAE

Name: Morishita Ryuichi Degree: MD, PhD Birthday: 5/12/1962 Birthplace: Japan Citizenship: Japanese

Carrier:		
4/81-3/87	MD(3/87)	Osaka University Medical School, Osaka, Japan Medicine
4/87-3/91	PhD(3/91)	Osaka University Medical School, Osaka, Japan Medicine
4/91-8/91	Postdoctoral Fellow	Osaka University Medical School Department of Geriatric Medicine (T. Ogihara)
8/91-4/94	Postdoctoral Fellow	Stanford University School of Medicine, Division of Cardiovascular Medicine
5/94-96/9	Senior Research Associate	(Victor J. Dzau)
313 ( 3013	Selifor Research / Associa	Osaka University Medical School
5/94-96/8	Visiting Instructor	Department of Geriatric Medicine (T. Ogihara) Stanford University School of Medicine, Division of Cardiovascular Medicine
4/95-96/9	Research Fellow of the Is	(Victor J. Dzau)  apan Society for the Promotion of Science
10/96-10/98	Assistant Professor	Department of Geriatric Medicine (T. Ogihara)
10/50 10/50	11351544111 110105501	Osaka University Medical School
5/94-present	Chief	Section of Gene Therapy
<b>F</b>		Department of Geriatric Medicine (T. Ogihara)
		Osaka University Medical School
10/98-03/2004	Associate Professor	Department of Geriatric Medicine (T. Ogihara)
		Osaka University Medical School
10/98-03/2004	Associate Professor	Division of Gene Therapy Science (Y. Kaneda)
		Osaka University Medical School
10/98-03/2004	Chief	Section of Cardiovascular Medicine
		Division of Gene Therapy Science (Y. Kaneda)
		Osaka University Medical School
01/2000-present	Visiting Professor	The University of Hong Kong
03/2003-present	Professor	Division of Clinical Gene Therapy
		Osaka University Medical School
		(Donated by Dai-ichi Pharmaceutical)

Honors:			
1991	Award in Japan Research Foundation for Clinical Pharmacology		
1991-199	Award in Japan Heart Foundation		
1992-199	Postdoctoral Fellowship Grant in American Heart Association, California		
1993	Award in American Federation Clinical Research		
1993	Upjohn Young Investigator Award in Cardiology, Stanford University		
1994	Young Investigator Award of the Dr. C & F. Demuth Medical Foundation at the		
	15th Scientific Meeting of the International Society of hypertension in Melbourne,		
	Australia.		
1994	Award in American Federation Clinical Research		
1994	Award in Second Conference "Hypertension and Vascular Metabolisms"		
1994	Young Investigator Award in Japan Vascular Disease Research Foundation		
1996	Young Investigator Award (First winner) in Japanese Circulation Society		
1996	Young Investigator Award in Japanese Atherosclerosis Society		
1996	Harry Goldbratt Award in Council of High Blood Pressure, American Heart		
	Association		
1997	Young Investigator Award in 11th International Symposium on Atherosclerosis		
1997	Young Investigator Award in 1st annual meeting of the Society of Cardiovascular		
	& Endocrinology		
1998	Young Investigator Award in 71th annual meeting of the Society of		
	Endocrinology		
1999	Young Investigator Award in the annual meeting of the Society of Pharmacology		
1999	Award in Japanese of Japan Medical Society		
2001	Takamine Jokichi Award in 5th annual meeting of the Society of Cardiovascular		
	Endocrinology		
2001	Young Investigator Award in 2nd annual meeting of the Japanese Society of Hypertension		
2002	Sato Award in 27 <sup>th</sup> annual meeting of the Japanese Circulation Society		
2003	Japan Innovator Award in 1st meeting (Nikkei BP)		
2003	Award from Minister of Education in 1 <sup>st</sup> meeting		
2005	Invitrogen-Nature-Biotechnology Award		
_000	and the second discontinuous d		

Editorial: Circulation Research (1998-), Circulation (1999-), Hypertension (2006-), Journal of Atherosclerosis and Thrombosis (Associate Editor; 1999-2004), Japanese Circulation Journal (2000-), Current Drug Targets (2000-), Current Gene Therapy (2000-), Heart (2000-), Journal of Hypertension (2000-2003), Expert Review of Cardiovascular Therapy (2002-), Cancer Therapy (2003-), Medicinal Chemistry Reviews (2003-), Gene Therapy & Molecular Biology (2004-), Current Hypertension Reviews (2004-), Current Cardiology Reviews (2004-), Geriatrics Gerontology International (2004-), Expert Opinion on Therapeutic Targets (2004-), Recent Patent Reviews On Cardiovascular Drug Discovery (2005-), Current Medicinal Chemistry (2006-), Recent Patents in Biotechnology (2006-), International Journal of Biomedical Science (2006-)

#### **Professional Societies:**

Board; Japanese Society of Vascular Biology Organization, Society for Intellectual Property

Fellow: Council of Circulation, American Heart Association, Council of Atherosclerosis, American Heart Association, Council of High Blood Pressure, American Heart Association, American College of Angiology, Japanese Society of Nephrology, Japanese Society of Pharmacology, Japanese Society of Atherosclerosis, Japanese Society of Endocrinology, Japanese Society of Cardiovascular & Endocrinology

Member, Council of Basic Science, American Heart Association, International Society of

Nephrology, American Society of Gene Therapy, North American Vascular Biology Organization, Japanese Society of Hypertension, Japanese Society of Circulation, Japanese Society of Gerontology, Japanese Society of Internal Medicine Japanese Society of Gene Therapy

#### **BIBLIOGRAPHY**

#### Original articles

- 1. Higaki J, Ogihara T, Nakamaru M, <u>Morishita R</u>, Kumahara Y. Effects of carvedilol on plasma hormonal and biochemical factors and renal function in Japanese patients with essential hypertension. Drugs 1988,36:64-68.
- 2. Higaki J; Okuno M; Nakamaru M; <u>Morishita R</u>; Himeno S; Ogihara T. Changes in plasma active renin and prorenin after endoscopic retrograde pancreatography. Methods and Findings in Experimental and Clinical Pharmacology 1989;11:703-705.
- 3. <u>Morishita R</u>; Higaki J; Ogihara T. Endothelin stimulates aldosterone biosynthesis by dispersed rabbit adreno-capsular cells. Biochemical and Biophysical Research Communications 1989;160:628-632.
- 4. Higaki J; <u>Morishita R</u>; Ogihara T. Measurements of enzyme activity and immunoreactivity of plasma renin. Limitations of the conventional method for assay of plasma renin catalytic activity. Journal of Clinical Chemistry and Clinical Biochemistry, 1989;27:201-204.
- 5. Higaki J; Morishita R; Ogihara T; Nishiura M. Effects of aging on human plasma renin: simultaneous multiple assays of enzyme activity and immunoactivity of plasma renin. Acta Endocrinologica, 1989;120:81-86.
- 6. Higaki J, <u>Morishita R</u>, Nakamaru M, Mikami H, Ogihara T. Thiazide-induced cerebral infarction in an elderly hypertensive patients with carotid artery stenosis. Advances in Therapy. 1989;6:180-185.
- 7. Morishita R; Higaki J; Mikami H; Ogihara T. Effect of cyclosporin A on aldosterone production by dispersed rabbit adreno-capsular cells. Life Sciences, 1990, 46:1985-1989.
- 8. Higaki J, Mikami H, Otsuka A, Iwatsubo H, Katahira K, Moriguchi A, Morishita R, Hata T, Ogihara T. Effect of two years of enalapril treatment on the quality of ligfe of elderly patients with essential hypertension complicated by other diseases. Current Therapeutic Research 1990;47:620-626.
- 9. Nakamura F; Nagano M; Higaki J; Higashimori K; Katahira K; Morishita R; Mikami H; Ogihara T. Regression of left ventricular hypertrophy by angiotensin converting enzyme inhibitor in reduced renal mass hypertensive rats. Journal of Hypertension. 1991;9:S398-399.
- 10. Morishita R; Higaki J; Nagano M; Mikami H; Ogihara T; Tanaka T; Ishii K; Okunishi H; Miyazaki M. Consistent activation of prorenin mRNA in renal hypertensive rats. Canadian Journal of Physiology and Pharmacology, 1991;69:1364-1366.
- 11. Higaki J; Miya A; Miki T; Morishita R; Mikami H; Takai S; Ogihara T. Contribution of the activation of the ras oncogene to the evolution of aldosterone- and renin-secreting tumors. Journal of Hypertension 1991;9:135-137.
- 12. Morishita R; Higaki J; Okunishi H; Tanaka T; Ishii K; Nagano M; Mikami H; Ogihara T; Murakami K; Miyazaki M. Changes in gene expression of the renin-angiotensin system in two-kidney, one clip hypertensive rats. Journal of Hypertension, 1991;9:187-192.
- 13. Morishita R, Higaki J, Okunishi H, Kawamoto T, Ishii K, Nakamura F, Katahira K, Nagano M, Mikami H, Miyazaki M, Ogihara T. Effect of long term treatment with an angiotensin converting enzyme inhibitor on the renin angiotensin system in spontaneously hypertensive rats. Clinical and Experimental Pharmacology and Physiology 1991;18:685-690.
- 14. <u>Morishita R.</u> Higaki J. Nakamura F. Tomita N. Yu H, Nagano M, Mikami H, Ogihara T. Role of vascular angiotensin II in hypertension induced vascular hypertrophy of spontaneously hypertensive rats. Blood Pressure 1992;1:41-47.
- 15. <u>Morishita R</u>; Higaki J; Katahira K, Nakamura F, Nagano M; Mikami H; Okunishi H, Miyazaki M, Ogihara T. Investigation of the fectors of the renin angiotensin system in renal hypertensive rats. Journal of Physiology 1992;446:370.
- 16. Katsuya T, Higaki J, Miki T, Nakura J, Ikegami H, Morishita R, Nagano M, Higashimori K, Nakamura F, Mikami H, Ogihara T. Substrain comparison of genetically hypertensive rats using DNA fingerprinting, and genetic analysis of blood pressure in the inbred rats. The

- Tohoku Journal of Experimental Medicine 1992;165:253-260.
- 17. Ogihara T, Ishikawa H, Mikami H, Nonaka H, Higaki J, Kagoshima T, Otsuka A, Higashimori K, Morishita R, Ishikawa K. Effects of long-term treatment with delapril on quality of life in hypertensive patients. Current Therapeutic Research 1992;51:254-260.
- 18. Ogihara T, Mikami H, Otsuka A, Nagano M, Matsuo K, Saeki S, <u>Morishita R</u>, Suzuki T, Naka T, Hata T. Effect of nilvadipine on quality of life in patients with essential hypertension. Current Therapeutic Research 1992;51:154-162.
- 19. Ogihara T, Nakagawa M, Ishikawa H, Mikami H, Takeda K, Nonaka H, Higaki J, Saeki S, Kagoshima T, Matsuo K, <u>Morishita R</u>, Ishikawa K. The effects of temocapril, a new angiotensin converting enzyme inhibitor, on the quality of life in hypertensive patients. Current Therapeutics Research 1992;52:81-88.
- 20. Tomita N, Higaki J, <u>Morishita R</u>, Kato K, Mikami H, Kaneda Y, Ogihara T. Direct in vivo gene introduction into rat kidney. Biochemical and Biophysical Research Communications 1992;186:129-34.
- 21. <u>Morishita R</u>, Higaki J, Nakamura F, Tomita N, Yu H, Nagano M, Mikami H, Ogihara T. Regression of hypertension-induced vascular hypertrophy by an ACE inhibitor and calcium antagonist in the spontaneously hypertensive rat. Blood Pressure 1992;3:41-47.
- 22. Morishita R; Higaki J; Miyazaki M; Ogihara T. Possible role of the vascular renin-angiotensin system in hypertension and vascular hypertrophy. Hypertension, 1992 19(Suppl):II62-7.
- 23. <u>Morishita R</u>; Higaki J; Mikami H; Matsukawa N; Ogihara T. Effect of aging on the renin angiotensin system in patients with renovascular hypertension. Methods and Findings in Experimental and Clinical Pharmacology, 1992;14:17-21.
- 24. Dzau VJ. Morishita R, Gibbons GH. Gene therapy in the cardiovascular diseases. Trends in Biotechnology 1993;11:205-210.
- 25. Morishita R, Gibbons GH, Kaneda Y, Ogihara T, Dzau VJ. Novel in vitro gene transfer method for study of local modulators in vascular smooth muscle cells. Hypertension 1993;21:894-899.
- 26. <u>Morishita R</u>; Higaki J; Okunishi H, Nakamura F, Nagano M; Mikami H; Ishii K, Miyazaki M, Ogihara T. Role of tissue renin angiotensin system in two-kidney, one clip hypertensive rats. American Journal of Physiology 1993;264:F510-514.
- 27. Nakamura F, Nagano M, Higaki J, HIgashimori K, <u>Morishita R</u>, Mikami H, Ogihara T. The angiotensin-converting enzyme inhibitor, perindopril, prevents cardiac hypertrophy in low-renin hypertensive rats. Clinical and Experimental Pharmacology and Physiology 1993;20:135-140.
- 28. Morishita R. Higaki J. Nagano M. Nakamura F. Tomita N. Zhao Y. Mikami H, Miyazaki M, Ogihara T. Discrepancy between renin mRNA and plasma renin level in angiotensin-converting enzyme inhibitor-treated rats. Clinical and Experimental Pharmacology and Physiology 1993;20:15-20.
- 29. <u>Morishita R</u>, Gibbons GH, Kaneda Y, Ogihara T, Dzau VJ. Novel and effective gene transfer technique for study of vascular renin angiotensin system. Journal of Clinical Investigation 1993;91:2580-2585
- 30. Morishita R, Gibbons GH, Ellison KE, Nakajima M, Zhang L, Kaneda Y, Ogihara T, Dzau VJ. Single intraluminal delivery of antisense cdc 2 kinase and PCNA oligonucleotides results in chronic inhibition of neointimal hyperplasia. Proc Natl Acad Sci USA 1993;90:8474-8478.
- 31. Morishita R, Gibbons GH, Ellison KE, Nakajima M, Leyen HVL, Zhang L, Kaneda Y, Ogihara T, Dzau VJ. Antisense oligonucleotides directed at cell cycle regulatory genes as a target for restenosis therapy. Transactions of Association of American Physicians 1993;106:54-61.
- 32. Tomita N, Higaki J, Kaneda Y, Yu H, <u>Morishita R</u>, Mikami H, Ogihara T. Hypertensive rats produced by in vivo introduction of the human renin gene. Circulation Research 1993;73:898-905.

- 33. Yu H, Rakugi H, Higaki J, <u>Morishita R</u>, Mikami H, Ogihara T. The role of activated vascular angiotensin II generation in vascular hypertrophy in one-kidney, one clip hypertensive rats. Journal of Hypertension 1993;11:1347-1355.
- 34. Morishita R, Gibbons GH, Ellison KE, Nakajima M, Leyen HVL, Zhang L, Kaneda Y, Ogihara T, Dzau VJ. Intimal hyperplasia after vascular injury is inhibited by antisense cdk 2 kinase oligonucleotides. Journal of Clinical Investigation 1994;93:1458-1464.
- 35. Morishita R, Rakugi H, Higaki J, Tomita N, Nakamura F, Yu H, Katsuya T, Mikami H, Ogihara T. Differential regulation of brain angiotensin II in genetically hypertensive and normotensive rats after nephrectomy. Blood Pressure 1994:3;265-269.
- 36. Morishita R, Gibbons GH, Pratt E, Tomita N, Kaneda Y, Ogihara T- Dzau VJ. Autocrine and paracrine effects of atrial natriuretic peptide gene transfer on vascular smooth muscle and endothelial cellular growth. Journal of Clinical Investigation 1994;94:824-829.
- 37. <u>Morishita R</u>, Gibbons GH, Ellison KE, Lee W, Zhang L, Kaneda Y, Ogihara T, Dzau VJ. Evidence for direct local effect of angiotensin in vascular hypertrophy: in vivo gene transfer of angiotensin converting enzyme. Journal of Clinical Investigation 94:978-984,1994.
- 38. Morishita R, Higaki J, Yu H, Tomita N, Nakamura F, Mikami H, Ogihara T. Effect of antihypertensive drugs on vascular hypertrophy and vascular angiotensin II in two-kidney, one clip hypertensive rats. Journal of Vascular Medicine and Biology 1994;5:38-45.
- 39. Morishita R., Gibbons GH, Kaneda Y, Ogihara T, Dzau VJ. Pharmacokinetics of antisense oligonucleotides (cyclin B1 and cdc 2 kinase) in the vessel wall: enhanced therapeutic utility for restenosis by HVJ-liposome method. Gene 1994;149:13-19.
- 40. Katahira K, Mikami H, Otuka A, Moriguchi A, Kohara K, Higashimori K, Okuda N, Nagano M, Morishita R, Ogihara T. Differential control of vascular tone and heart rate by different amino acid neurotransmitters in the rostral ventrolateral medulla of the rat. Clinical and Experimental Physiology and Pharamcology. 1994;21:545-556.
- 41. Tomita N, Higaki J, <u>Morishita R</u>, Kaneda Y, Ogihara T. A novel gene transfer technique mediated by HVJ (Sendai virus), nuclear protein and liposomes. Cancer Detection and Prevention. 1994;18:485-491.
- 42. <u>Morishita R</u>, Higaki J, Rakugi H, Tomita N, Yu H, Nakamura F, Mikami H, Ogihara T. Effect of antihypertensive drug on brain angiotensin II level in renal and spontaneously hypertensive rats. Clinical and Experimental Physiology and Pharmacology 1995;22;665-669.
- 43. Leyen H.V.L., Gibbons G.H., <u>Morishita R.</u>, Lewis N.P., Zhang L., Nakajima M., Kaneda Y., Cooke J.P., Dzau V.J. Gene therapy inhibiting neointimal vascular lesion: in vivo transfer of ec-nitric oxide synthase gene. Proc Natl Acad Sci USA 1995;92:1137-1141.
- 44. Morishita R, Gibbons GH, Horiuchi M, Ellison KE, Nakajima M, Zhang L, Kaneda Y, Ogihara T, Dzau VJ. A gene therapy strategy using a transcription factor decoy of the E2F binding site inhibits smooth muscle proliferation in vivo. Proc Natl Acad Sci 1995;92:5855-5859.
- 45. Tomita N, Morishita R, Higaki J, Aoki M, Nakamura Y, Mikami H, Fukamizu A, Murakami K, Kaneda Y, Ogihara T. Transient decrease in high blood pressure by in vivo transfer of antisense oligodeoxynucleotides against rat angiotensinogen. Hypertension 1995;26:131-136.
- 46. Yamada T, Horiuchi M, <u>Morishita R</u>, Zhang L, Pratt RE, Dzau VJ. In vivo identification of a negative regulatory element in the mouse renin gene using direct gene transfer. Journal of Clinical Investigation 1995;96:1230-1237.
- 47. Tomita N, Morishita R, Higaki J, Tomita S, Aoki M, Kaneda Y, Ogihara T. Role of angiotensinogen in blood pressure regulation in normotensive rats: application of a "loss of function" approach. Journal of Hypertension 1995;13;1767-1774.
- 48. Yamada K, Moriguchi A, Mikami H, Higaki J, Morishita R, Aoki M, Okuda N, Ogihara T. Importance of central amino acid neurotransmitter in the blood pressure regulation in the spontaneously hypertensive rat. Journal of Hypertension 1995;13;1624-1630.
- 49. Nakamura Y, Morishita R, Higaki J, Kida I, Aoki M, Moriguchi A, Yamada K, Hayashi S, Yo

- Y, Matsumoto K, Nakamura T. Expression of local hepatocyte growth factor system in vascular tissues. Biochemical Biophysics Research Communication 1995;215:483-488.
- 50. Nakajima M, Hutchinson H, Fujinaga M, Hayashida W, Morishita R, Zhang L, Horiuchi M, Pratt RE, Dzau VJ. The AT2 receptor antagonizes the growth effects of the AT1 receptor: gain of function study using gene transfer. Proceeding of National Academy of Sciences USA 1995;92:10663-10667.
- 51. Hara S, Morishita R, Tone Y, Yokoyama C, Inoue H, Kaneda Y, Ogihara T, Tanabe T. Overexpression of prostacyclin synthase inhibits growth of vascular smooth muscle cells. Biochemical Biophysics Research Communication 1995;216:862-867.
- 52. Sawa Y, Suzuki K, Bai HZ, Shirakura R, Morishita R, Kaneda Y, Matsuda H. Efficiency of in vivo gene transfection into transplanted rat heart by coronary infusion of HVJ-liposome. Circulation 1995;92:II-479-482.
- 53. Ellison KE, Bishopric NH, Webster KA, <u>Morishita R</u>, Gibbson GH, Kaneda Y, Sato B, Dzau VJ, Fusigenic liposome-mediated DNA transfer into cardiac myocytes. Journal of Molecular and Cellular Cardiology 1995;28:1385-1399.
- 54. Morishita R, Higaki J, Tomita N, Aoki M, Moriguchi A, Tamura K, Murakami K, Kaneda Y, Ogihara T. Role of transcriptional cis-elements, angiotensinogen gene-activating element, of angiotensinogen gene in blood pressure regulation. Hypertension 1996;27:502-507.
- 55. Hayashi S, Morishita R, Higaki J, Aoki M, Moriguchi A, Kida I, Sawa Y, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Autocrine-paracrine effects of over-expression of hepatocyte growth factor gene on growth of endothelial cells. Biochemical Biophysics Research Communication 1996;220:539-545.
- 56. Tomita N, Morishita R, Higaki J, Tomita S, Aoki M, Ogihara T, Kaneda Y. *In vivo* gene transfer of insulin gene into neonatal rats by HVJ-liposome method resulted in sustained transgene expression. Gene Therapy 1996:3;477-482.
- 57. Tomita N, Morishita R, Higaki J, Tomita S, Aoki M, Kaneda Y, Ogihara T. Transfection of renin gene into neonates results in sustained gene expression, but fails to produce sustained hypertension. Biochemical Biophysics Research Communication 1996;224:43-49.
- 58. Nakamura Y, Morishita R, Higaki J, Kida I, Aoki M, Moriguchi A, Yamada K, Hayashi S, Yo Y, Nakano H, Matsumoto K, Nakamura T, Ogihara T. Hepatocyte growth factor (HGF) ia a novel member of endothelium-specific growth factors: additive stimulatory effect of HGF with basic fibroblast growth factor, but not vascular endothelial growth factor. Journal of Hypertension 1996;14:1067-1072.
- 59. Yamada K, Moriguchi A, Morishita R, Aoki M, Nakamura Y, Mikami H, Oshima T, Ninomiya M, Kaneda Y, Higaki J, Ogihara T. Efficient oligonucleotides delivery using HVJ-liposome method in the central nervous system. American Journal of Physiology 1996;40:R1212-1220.
- 60. Nakamura Y, <u>Morishita R</u>, Nakamura S, Aoki M, Moriguchi A, Matsumoto K, Nakamura T, Higaki J, Ogihara T. A vascular modulator, hepatocyte growth factor, is associated with systolic pressure. Hypertension 1996;28:409-413
- 61. Hayashi S, <u>Morishita R</u>, Aoki M, Moriguchi A, Kida I, Nakajima M, Kaneda Y, Higaki J, Ogihara T. <u>In vivo</u> transfer of gene and oligodeoxynucleotides into skin of fetal rats by incubation in amniotic fluid. Gene Therapy 1996;3:878-885.
- 62. Yonemitsu Y, Kaneda Y, <u>Morishita R</u>, Nakagawa K, Nakashima Y, Sueishi K. Characterization of in vivo gene transfer into the arterial wall mediated by the Sendai virus (Hemagglutinating Virus of Japan) liposomes: an effective tool for the in vivo study of arterial diseases. Laboratory Investigation 1996;75:313-323.
- 63. Morishita R, Gibbons GH, Horiuchi M, Nakajima M, Ellison KE, Lee W, Kaneda Y, Ogihara T, Dzau VJ, Molecular delivery system for antisense olgonucleotides: enhanced effectiveness of antisense oligonucleotides by HVJ-liposome mediated transfer. Journal of Cardiovascular Pharmacology & Therapeutics 1996;2:213-222.

- 64. Morishita R, Nakamura S, Nakamura Y, Aoki M, Moriguchi A, Kida I, Yo Y, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Potential role of endothelium-specific growth factor, hepatocyte growth factor, on endothelial damage in diabetes mellitus. Diabetes 1997;46:138-142.
- 65. Aoki M, Morishita R, Higaki J, Moriguchi A, Hayashi S, Matsushita H, Kida I, Tomita N, Sawa Y, Kaneda Y, Ogihara T. Survival of grafts of genetically modified cardiac myocytes transfected with FITC-labeled oligodeoxynucleotides and β-galactosidase gene in non-infarcted area, but not myocardial infarcted area. Gene Therapy 1997;4:120-127.
- 66. Aoki M, <u>Morishita R</u>, Higaki J, Moriguchi A, Kida I, Hayashi S, Matsushita H, Kaneda Y, Ogihara T. <u>In vivo</u> transfer efficiency of antisense oligonucleotides into the myocardium using HVJ-liposome method. Biochemical Biophysics Research Communications 1997;231:540-545.
- 67. Aoki M, Morishita R, Muraishi A, Moriguchi A, Sugimoto T, Maeda K, Dzau VJ, Kaneda Y, Higaki J, Ogihara T. Efficient in vivo gene transfer into heart in rat myocardial infarction model using HVJ (Hemagglutinating Virus of Japan)-liposome method. Journal of Molecular and Cellular Cardiology 1997;29:949-959.
- 68. Rakugi H, Okamura A, Kamide K, Ohishi M, Sasamura H, <u>Morishita R</u>, Higaki J, Ogihara T. Recognition of tissue- and subtype-specific modulation of angiotensin II receptors using antibodies against AT1 and AT2 receptors. Hypertension Research 1997;20:51-55.
- 69. Morishita R, Higaki J, Hayashi S, Yo Y, Aoki M, Nakamura S, Moriguchi A, Matsushita H, Matsumoto K, Nakamura T, Ogihara T. Role of hepatocyte growth factor in endothelial regulation: Prevention of high D-glucose-induced endothelial cell death by prostaglandins and phosphodiesterase type 3 inhibitor. Diabetologia 1997;40:1053-1061.
- 70. Tomita T, Hashimoto H, Tomita N, Morishita R, Bak Lee S, Hayashida K, Nakamura N, Yonenobu K, Kaneda Y, Ochi T. In vivo eirect gene transfer into articular cartilage by intraarticular injection mediated by HVJ (Sendai virus) and liposomes. Arthritis & Rheumatism 1997;40:901-906.
- 71. Morishita R, Sugimoto T, Aoki M, Kida I, Tomita N, Moriguchi A, Maeda K, Sawa Y, Kaneda Y, Higaki J, Ogihara T. In Vivo Transfection of cis element "decoy" against NFkB binding site prevented myocardial infarction as gene therapy. Nature Medicine 1997;3:894-899.
- 72. Suzuki J, Isobe M, Morishita R, Aoki M, Horie S, Okubo Y, Kaneda Y, Sawa Y, Matsuda H, Ogihara T, Sekiguchi M. Prevention of graft coronary arteriosclerosis by antisense cdk 2 kinase oligonucleotide. Nature Medicine 1997;3:900-903.
- 73. Aoki M, Morishita R, Matsushita H, Nakano N, Hayashi S, Tomita N, Yamamoto K, Moriguchi A, Higaki J, Ogihara T. Serum deprivation induced apoptosis accompanied by up-regulation of p53 and bax in human aortic vascular smooth muscle cells. Heart & Vessels 1997;S12:71-75.
- 74. Sawa Y, Morishita R, Suzuki K, Kagisaki K, Kaneda Y, Maeda K, Kadoba K, Matsuda H. A novel strategy for myocardial protection using in vivo transfection of cis element "decoy" against NFkB binding site: evidence for a role of NFkB in ischemic-reperfusion injury. Circulation 1997;96:II-280-285.
- 75. Nakano N, Moriguchi A, <u>Morishita R</u>, Kida I, Tomita N, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Role of angiotensin II in the regulation of a novel vascular modulator, hepatocyte growth factor, in experimental hypertensive rats. Hypertension 1997;30:1448-1454.
- 76. Yo Y, Morishita R, Yamamoto K, Tomita N, Kida I, Hayashi S, Moriguchi A, Kato S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Actions of Hepatocyte Growth Factor as a Local Modulator in the Kidney: Potential Role in Pathogenesis of Renal Disease. Kidney International 1998;53:50-58.
- 77. Matsushita H, Morishita R, Kida I, Aoki M, Hayashi S, Tomita N, Yamamoto K, Moriguchi A, Noda A, Kaneda Y, Higaki J, Ogihara T. Inhibition of growth of human vascular smooth muscle cells by over-expression of p21 gene through induction of apoptosis. Hypertension 1998;31:493-498.

- 78. Matsumoto T, Komori K, Yonemitsu Y, <u>Morishita R</u>, Sueishi K, Kaneda Y, Sugimachi K. Hemagglutinating virus of Japan-liposome mediated gene transfer of endothelial cell nitric oxide synthase inhibits intimal hyperplasia of canine vein grafts under conditions of poor runoff. Journal of Vascular Surgery 1998;27:125-144.
- 79. Nakamura S, Moriguchi A, <u>Morishita R</u>, Aoki M, Yo Y, Hayashi S, Nakano N, Katsuya T, Nakata S, Takami S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. A novel vascular modulator, hepatocyte growth factor (HGF), as a potential index of the severity of hypertension. Biochemical Biophysics Research Communication 1998;242:238-243
- 80. Morishita R, Gibbons GH, Horiuchi M, Kaneda Y, Ogihara T, Dzau VJ. Role of AP-1 complex in angiotensin II-mediated transforming growth factor-β expression and growth of smooth muscle cells:using decoy approach against AP-1 binding site. Biochemical Biophysics Research Communication 1998;243:361-367.
- 81. Hirano T, Fujimoto J, Ueki T, Yamamoto H, Iwasaki T, Morishita R, Sawa Y, Kaneda Y, Takahashi H, Okamoto E. Persistent gene expression in rat liver in vivo by repetitive transfections using HVJ-liposome. Gene Therapy 1998;5:459-464.
- 82. Ono S, Date I, Onoda K, Shiota T, Ohmoto T, Ninomiya Y, Asari S, <u>Morishita R</u>. Decoy administration of NFkB into the subarachnoid space for cerebral angiopathy. Human Gene Therapy 1998;9:1003-1011.
- 83. Suzuki J, Isobe M, Morishita R; Aoki M, Yamazaki S, Kaneda Y, Sawa Y, Matsuda H, Ogihara T, Horie S, Okubo Y, Sekiguchi M. Antisense Cdk2 kinase oligodeoxynucleotide inhibits ICAM-1 expression in murine cardiac allograft arteriopathy. Transplant Proc 1998;30:89-90.
- 84. Tomita N, Morishita R, Tomita S, Yamamoto K, Aoki M, Matsushita H, Hayashi S, Higaki J, Ogihara T. Transcription factor decoy for NFkB inhibits TNF-α-induced IL-6 and ICAM-1 expression in endothelial cells. Journal of Hypertension 1998;16:993-1000.
- 85. Isobe M, Suzuki J, Morishita R, Aoki M, Amano J, Kaneda Y, Sawa Y, Matsuda H, Ogihara T, Horie S, Okubo Y, Sekiguchi M. Downregulation of endothelin-1 expression in allograft coronary arteries after gene therapy targeting cdk2 kinase. Transplantation Proceedings 1998;30:1007-1008
- 86. Bai HZ, Morishita R, Kida I, Yamakawa T, Zhang WD, Aoki M, Matsushita H, Noda A, Nagai R, Kaneda Y, Higaki J, Ogihara T, Sawa Y, Matsuda H. Inhibition of intimal hyperplasia after vein grafting by in vivo transfer of human senescent cell-derived inhibitor-1 gene. Gene Therapy 1998;5:761-769.
- 87. Morishita R, Yamada S, Higaki J, Tomita N, Kida I, Aoki M, Moriguchi A, Hayashi S, Sakurabayashi I, Kaneda Y, Ogihara T. Conditioned medium from HepG2 cells transfected with human apolipoprotein (a) gene stimulates growth of human vascular smooth muscle cells: effects of over-expression of human apolipoprotein (a) gene. Hypertension 1998;32:215-222.
- 88. Morishita R, Yamamoto K, Yamada S, Matsushita H, Tomita N, Sakurabayashi I, Kaneda Y, Moriguchi A, Higaki J, Ogihara T. Stimulatory effect of lipoprotein (a) on proliferation of human mesangial cells: role of lipoprotein (a) in renal disease. Biochemical Biophysics Research Communication 1998;249:313-320
- 89. Nakano N, Morishita R, Moriguchi A, Nakamura Y, Hayashi S, Aoki M, Kida I, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Negative regulation of local hepatocyte growth factor (HGF) expression by angiotensin II and transforming growth factor-β in blood vessels: potential role of HGF in cardiovascular disease. Hypertension 1998;32:444-451
- 90. Yo Y, Morishita R, Nakamura S, Tomita N, Yamamoto K, Moriguchi A, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Potential role of hepatocyte growth factor in the maintenance of renal structure: anti-apoptotic action of HGF on epithelial cells. Kidney International 1998:54:1128-1138
- 91. Morishita R, Yamada S, Yamamoto K, Tomita N, Kida I, Sakurabayashi I, Kikuchi A, Kaneda Y, Lawn R, Higaki J, Ogihara T. Novel therapeutic strategy for atherosclerosis:

- ribozyme oligonucleotides against apolipoprotein (a) selectively inhibit apolipoprotein (a), but not plasminogen, gene expression. Circulation 1998;98:1898-1904
- 92. Sawa Y, Kaneda Y, Bai HZ, Suzuki K, Fujimoto J, <u>Morishita R</u>, Matsuda H. Efficient transfer of oligonucleotides and plasmid DNA into the whole heart through the coronary artery. Gene Therapy 1998;5;1472-1480.
- 93. Nakamura S, <u>Morishita R</u>, Moriguchi A, Yo Y, Nakamura Y, Hayashi S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Hepatocyte growth factor as a potential index of complication in diabetes mellitus. Journal of Hypertension 1998;16:2019-2026.
- 94. Kawamura I, <u>Morishita R</u>, Tomita N, Elizabeth L, Aketa M, Tsujimoto S, Manda T, Tomoi M, Kida I, Higakai J, Kaneda Y, Shimomura K, Ogihara T. Intratumoral injection of oligonucleotides to the NFkB binding site inhibits cachexia in a mouse tumor model. Gene Therapy 1999;6;91-97
- 95. Ueki T, Kaneda Y, Tsutsui H, Nakanishi K, Sawa Y, <u>Morishita R</u>, Matsumoto K, Nakamura T, Takahashi H, Oakamoto E, Fujimoto J. Curative gene therapy of liver cirrhosis by hepatocyte growth factor in rats. Nature Medicine 1999;5:226-230
- 96. Todaka T, Yokoyama C, Yanamoto H, Hashimoto N, Nagata I, Tsukahara T, Hara S, Hatae T, Morishita R, Aoki M, Ogihara T, Kaneda Y, Tanabe T. Gene transfer of human prostacyclin synthase prevents neointimal formation after carotid balloon injury in rats. Stroke 1999;30:419-426.
- 97. Tomita S, Tomita N, Yamada T, Xhang L, Kaneda Y, Morishita R, Ogihara T, Dzau VJ, Horiuchi M. Transcription factor decoy to syudy the molecular mechanism of negative regulation of renin gene expression in the liver in vivo. Circulation Research 1999;84:1059-1066.
- 98. Morishita R, Nakamura S, Hayashi S, Taniyama Y, Moriguchi A, Nagano T, Taiji M, Noguchi H, Takeshita S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Therapeutic angiogenesis induced by human recombinant hepatocyte growth factor in rabbit hind limb ischemia model as "cytokine supplement therapy" Hypertension 1999;33:1379-1384
- 99. Nishii T, Moriguchi A, <u>Morishita R</u>, Yamada K, Nakamura S, Tomita N, Kaneda Y, Fukamizu A, Mikami H, Higaki J, Ogihara T. Angiotensinogen gene activating elements regulate blood pressure in the brain. Circulation Research 1999;85:257-263.
- 100.Aoki M, Morishita R, Matsushita H, Hayashi S, Nakagami H, Yamamoto K, Moriguchi A, Kaneda Y, Higaki J, Ogihara T. Inhibition of the p53 tumor suppressor gene resulted in growth of human aortic vascular smooth muscle cells (VSMC): potential role of p53 in regulation of VSMC growth. Hypertension 1999;34:192-200
- 101.Matsumoto K, Morishita R, Moriguchi A, Tomita N, Yo Y, Nishii T, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Prevention of renal damage by angiotensin II blockade, accompanied by increased renal hepatocyte growth factor in experimental hypertensive rats. Hypertension 1999;34:279-284
- 102. Nakamura S, Moriguchi A, Morishita R, Yamada K, Nishii T, Tomita N, Oishi M, Kaneda Y, Higaki J, Ogihara T. Activation of the brain angiotensin system by in vivo human angiotensin converting enzyme gene transfer in rats. Hypertension 1999;34:302-308.
- 103.Zhou WZ, Kaneda Y, Huang SKS, <u>Morishita R</u>, Hoon DSB. Protective immunization against melanoma by gp100 DNA HVJ-liposome vaccine. Gene Therapy 1999;6:1768-1773
- 104. Yamada S, Inoue K, Morishita R, Ogihara T, Kubono K, Kubo N, Abe A, Sakurabayashi I. A new Lp(a) assay that is unaffected by apo(a) size polymorphism. Clinica Chimica Acta 1999;287:29-43.
- 105. Suzuki J, Isobe M, Morishita R, Izawa A, Yamazaki S, Okubo Y, Kaneda Y, Sekiguchi M. E2F decoy suppresses E-selectin expression in murine cardiac allograft arteriopathy. Transplant Proc 1999;31:2018-2019
- 106. Suzuki J, Izawa A, <u>Morishita R</u>, Kaneda Y, Sawa Y, Okubo Y, Ogihara T, Sekiguchi M, Isobe M. Antisense Cdc2 kinase oligonucleotide inhibits adhesion molecule expression in cardiac

- allograft arteriopathy. Transplant Proc 1999;31:955-956
- 107. Suzuki J, Izawa A, <u>Morishita R</u>, Kaneda Y, Sawa Y, Okubo Y, Ogihara T, Sekiguchi M, Isobe M. Prevention of cardiac allograft arteriopathy by antisense Cdc2 kinase oligonucleotide. Transplant Proc 1999;31:867-868
- 108.Hayashi S, Morishita R, Nakamura S, Yamamoto K, Moriguchi A, Nagano T, Taizi M, Noguchi H, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Potential role of hepatocyte growth factor, a novel angiogenic growth factor, in peripheral arterial disease: down-regulation of HGF in response to hypoxia in vascular cells. Circulation 1999;100:II301-II308.
- 109.Zhou WZ, Hoon DSB, Huang SKS, Fujii S, Hashimoto K, <u>Morishita R</u>, Kaneda Y. RNA melanoma vaccine: induction of anti-tumor immunity by human gp100 mRNA immunization. Human Gene Therapy 1999;10:2719-2724
- 110.Tomita T, Takeuchi E, Tomita N, <u>Morishita R</u>, Kaneko M, Yamamoto K, Nakase T, Seki H, Kato K, Kaneda Y, Ochi T. Suppressed severity of collagen-induced arthritis by in vivo transfection of nuclear factor kappaB decoy oligodeoxynucleotides as a gene therapy. Arthritis Rheum 1999;42:2532-2542
- 111.Hayashi Si, <u>Morishita R</u>, Matsushita H, Nakagami H, Taniyama Y, Nakamura T, Aoki M, Yamamoto K, Higaki J, Ogihara T. Cyclic AMP Inhibited Proliferation of Human Aortic Vascular Smooth Muscle Cells, Accompanied by Induction of p53 and p21. Hypertension 2000;35:237-243
- 112.Higaki J, Aoki M, <u>Morishita R</u>, Kida I, Taniyama Y, Tomita N, Yamamoto K, Moriguchi A, Kaneda Y, Ogihara T. <u>In vivo</u> evidence of importance of cardiac angiotensin converting enzyme in the pathogenesis of cardiac hypertrophy. Arteriosclerosis, Thrombosis & Vascular Biology 2000;20:428-434.
- 113.Suzuki J, Isobe M, <u>Morishita R</u>, Nishikawa T, Amano J, Kaneda Y. Antisense bcl-x oligonucleotide induces apoptosis and prevents arterial neointimal formation in murine cardiac allografts. Cardiovascular Research 2000;45:783-787.
- 114.Kaneda Y, Saeki Y, Nakabayashi M, Zhou WZ, Wataya-Kaneda M, <u>Morishita R.</u> Enhancement of transgene expression by cotransfection of oriP plasmid with EBNA-1 expression vector. Human Gene Therapy 2000;11:471-479.
- 115. Aoki M, Morishita R, Taniyama Y, Kida I, Moriguchi A, Matsumoto K, Nakamura T, Kaneda Y, Higaki J, Ogihara T. Angiogenesis induced by hepatocyte growth factor in non-infarcted myocardium and infarcted myocardium: up-regulation of essential transcription factor for angiogenesis, ets. Gene Therapy 2000;7:417-427
- 116.Suzuki K, Nakashima H, Sawa Y, <u>Morishita R</u>, Matsuda H, Kaneda Y. Reconstituted fusion liposomes for gene transfer in vitro and in vivo. Gene Therapy & Regulation. 2000;1:65-77.
- 117. Morishita R, Gibbons GH, Tomita N, Zhang L, Kaneda Y, Ogihara T, Dzau VJ. Antisense oligodeoxynucleotide inhibition of vascular angiotensin converting enzyme expression attenuates neointimal formation: evidence for tissue ACE function. Arteriosclerosis, Thrombosis & Vascular Biology 2000;20:915-922.
- 118.Matsushita H, Morishita R, Aoki M, Tomita N, Taniyama Y, Nakagami H, Shimozato T, Higaki J, Kaneda Y, Ogihara T. Transfection of antisense p53 tumor suppressor gene oligodeoxynucleotides into rat carotid artery resulted in abnormal growth of vascular smooth muscle cells. Circulation 2000;101:1447-1452
- 119.Matsushita H, <u>Morishita R</u>, Nata T, Aoki M, Nakagami H, Taniyama Y, Yamamoto K, Higaki J, Kaneda Y, Ogihara T. Hypoxia induced endothelial apoptosis through NFkB-mediated bcl-2 suppression: <u>in vivo</u> evidence of importance of NFkB in endothelial cell regulation. Circulation Research 2000;86:974-981
- 120. Yamano H, Ura K, <u>Morishita R</u>, Nakajima H, Monden M, Kaneda Y. Amplification of transgene expression in vitro and in vivo using a novel inhibitor of histone deacteylase. Molecular Therapy 2000;1:574-580.
- 121. Taniyama Y, Morishita R, Nakagami H, Moriguchi A, Sakonjo H, Kim S, Matsumoto K,

- Nakamura T, Higaki J, Ogihara T. Potential contribution of a novel anti-fibrotic factor, hepatocyte growth factor, to prevention of myocardial fibrosis by angiotensin II blockade in cardiomyopathic hamster. Circulation 2000;102:246-252
- 122. Tomita N, Morishita R, Lan HY, Yamamoto K, Hashizume M, Notakae M, Toyosawa K, Fujitani B, Mu W, Nikolic-Paterson DJ, Atkins RC, Kaneda Y, Higaki J, Ogihara T. *In vivo* administration of a nlear transcription factor-kappaB decoy suppresses experimental crescentic glomerulonephritis. Journal of American Society of Nephrology 2000;11:1244-1252
- 123. Morishita R, Gibbons GH, Kaneda Y, Ogihara T, Dzau VJ. Systemic administration of HVJ viral coat-liposome complex containing human insulin vector decreases glucose level in diabetic mouse: a model of gene therapy. Biochemical Biophysics Research Communication 2000;273:666-674
- 124.Tomita T, Takano H, Tomita N, <u>Morishita R</u>, Kaneko M, Shi K, Takahi K, Nakase T, Kaneda Y, Yoshikawa H, Ochi T. Transcription factor decoy for nuclear factor kappaB inhibits cytokine and adhesion molecule expressions in synovial cells derived from rheumatoid arthritis. Rheumatology 2000;39:749-757
- 125.Tomita N, Morishita R, Tomita S, Gibbons GH, Zhang L, Horiuchi M, Kaneda Y, Higaki J, Ogihara T, Dzau VJ. Transcription factor decoy for NFkB inhibits TNF-α-induced cytokine and adhesion molecule expression in vivo. Gene Therapy 2000;7:1326-1332
- 126. Yamamoto K, Morishita R, Tomita N, Shimozato T, Nakagami H, Kikuchi A, Aoki M, Higaki J, Kaneda Y, Ogihara T. Ribozyme oligonucleotides against transforming growth factor-β inhibited neointimal formation after vascular injury in rat model: potential application of ribozyme strategy to treat cardiovascular disease. Circulation 2000;102:1308-1314
- 127.Kitamoto S, Egashira K, Kataoka C, Koyanagi M, Katoh M, Shimokawa H, Morishita R, Kaneda Y, Sueishi K, Takeshita A. Increased activity of nuclear factor-kappaB participates in cardiovascular remodeling induced by chronic inhibition of nitric oxide synthesis in rats. Circulation. 2000;102:806-812.
- 128.Egashira K, Koyanagi M, Kitamoto S, Ni W, Kataoka C, <u>Morishita R</u>, Kaneda Y, Akiyama C, Nishida K, Takeshita A. Anti-monocyte chemoattractant protein-1 gene therapy inhibits vascular remodeling in rats. FASEB Journal 2000;14:1974-1978
- 129.Suzuki J, Isobe M, <u>Morishita R</u>, Nishikawa T, Amano J, Kaneda Y. Prevention of cardiac allograft arteriosclerosis using antisense proliferating-cell nuclear antigen oligonucleotide. Transplantation 2000;70:398-400
- 130. Morishita R. Adventure of gene therapy into the brain: a new era for cardiovascular gene therapy (Editorial). Circulation research 2000;87:719-721.
- 131.Nakagami H, Morishita R, Yamamoto K, Taniyama Y, Aoki M, Kim S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Anti-apoptotic action of hepatocyte growth factor (HGF) through mitogen-activated protein kinase on human aortic endothelial cells. Journal of Hypertension 2000;18:1411-1420
- 132.Hayashi K, Nakamura S, <u>Morishita R</u>, Moriguchi A, Aoki M, Matsumoto K, Nakamura T, Kaneda Y, Sakai N, Ogihara T. In vivo transfer of human hepatocyte growth factor gene accelerates re-endothelialization and inhibits neointimal formation after balloon injury in rat model. Gene Therapy 2000;7:1664-1671
- 133. Suzuki J, Morishita R, Amano J, Kaneda Y, Isobe M. Decoy against nuclear factor-kappa B attenuates myocardial cell infiltration and arterial neointimal formation in murine cardiac allografts. Gene Therapy 2000;7:1847-1852.
- 134. Yamada S, <u>Morishita R</u>, Nakamura S, Ogihara T, Kusumi Y, Sakurai I, Kubo N, Sakurabayashi I. Development of antibody against epitope of lipoprotein (a) modified by oxidation: evaluation of new enzyme-linked immunoassay for oxidized Lp (a). Circulation 2000;102:1639-1644.
- 135.Nagaya N, Yokoyama C, Kyotani S, Shimonishi M, <u>Morishita R</u>, Uematsu M, Nishikimi T, Nakanishi N, Ogihara T, Yamagishi M, Miyatake K, Kaneda Y, Tanabe T. Gene transfer of

- human prostacyclin synthase ameliorates monocrotaline-induced pulmonary hypertension in rats. Circulation 2000;102:2005-2010.
- 136.Hayashi I, Majima M, Fujita T, Okumura T, Kumagai Y, Tomita N, <u>Morishita R</u>, Higaki J, Ogihara T. In vivo transfer of antisense oligonucleotide against urinary kininase blunts deoxycorticosterone acetate-salt hypertension in rats. Br J Pharmacol. 2000;131:820-826.
- 137.Kawauchi M, Suzuki J, <u>Morishita R</u>, Wada Y, Izawa A, Tomita N, Amano J, Kaneda Y, Ogihara T, Takamoto S, Isobe M. Gene Therapy for attenuating cardiac allograft arteriopathy using ex vivo E2F decoy transfection by HVJ-AVE-liposome method in mice and nonhuman primates. Circulation Research 2000;87:1063-1068
- 138.Okada M, Hojo Y, Ikeda U, Takahashi M, Takizawa T, <u>Morishita R</u>, Shimada K. Interaction between monocytes and vascular smooth muscle cells induces expression of hepatocyte growth factor. J Hypertens. 2000;18:1825-1831.
- 139.Shinmura K, Morishita R, Aoki M, Higaki J, Ogihara T, Kaneda Y, Tani M. Catheter-delivered in vivo gene transfer into rat myocardium using the fusigenic liposomal mediated method. Jpn Heart J. 2000;41:633-647.
- 140. Sugimoto T, Mine H, Horii Y, Takahashi K, Nagai R, Morishita R, Komada M, Asada Y, Sawada T. Neuroblastoma cell lines showing smooth muscle cell phenotypes. Diagn Mol Pathol. 2000;9:221-228.
- 141. Higaki J, Katsuya T, Morishita R, Ogihara T. Hypertension and Genes. Internal Medicine. 2001;40:144-147.
- 142. Nakagami H, <u>Morishita R</u>, Yamamoto K, Taniyama Y, Aoki M, Matsumoto K, Nakamura T, Kaneda Y, Horiuchi M, Ogihara T. Mitogenic and anti-apoptotic actions of HGF through ERK, STAT3 and Akt in endothelial cells. Hypertension 2001;37:581-586
- 143.Matsumoto K, <u>Morishita R</u>, Moriguchi A, Tomita N, Aoki M, Sakonjo H, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Inhibition of neointima by ACE inhibitor in porcine coronary artery balloon injury model. Hypertension 2001;37:270-274
- 144.Tomita N, Morishita R, Higaki J, Ogihara T. Novel molecular therapeutic approach to cardiovascular disease based on hepatocyte growth factor. Journal of Atherosclerosis and Thrombosis. 2001;7:1-7.
- 145. Aoki M, <u>Morishita R</u>, Taniyama Y, Kaneda Y, Ogihara T. Therapeutic angiogenesis induced by hepatocyte growth factor: potential gene therapy for ischemic diseases. Journal of Atherosclerosis and Thrombosis. 2001;7:71-76.
- 146. Taniyama Y, Morishita R, Aoki M, Nakagami H, Yamamoto K, Yamazaki K, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Therapeutic angiogenesis induced by human hepatocyte growth factor gene in rat and rabbit hind limb ischemia models: preclinical study for treatment of peripheral arterial disease. Gene Therapy 2001;8:181-189
- 147. Tsuboniwa N, Morishita R, Hirano T, Fujimoto J, Furukawa S, Kikumori M, Okuyama A, Kaneda Y. Safety evaluation of hemagglutinating virus of japan-artificial viral envelope liposomes in nonhuman primates. Hum Gene Ther. 2001;12:469-487.
- 148.Kawauchi M, Suzuki J, Wada Y, <u>Morishita R</u>, Kaneda Y, Isobe M, Amano J, Takamoto S. Downregulation of nuclear factor kappa B expression in primate cardiac allograft arteries after E2F decoy transfection. Transplant Proc. 2001;33:451.
- 149. Tomita N, Morishita R, Tomita S, Kaneda Y, Higaki J, Ogihara T, Horiuchi M. Inhibition of TNF-alpha, induced cytokine and adhesion molecule. expression in glomerular cells in vitro and in vivo by transcription factor decoy for NFkappaB. Experimental Nephrology. 2001;9:181-190
- 150. Yamamoto K, Morishita R, Hayashi S, Matsushita H, Nakagami H, Moriguchi A, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Contribution of Bcl-2. but not Bcl-xL and Bax, to anti-apoptotic actions of hepatocyte growth factor in hypoxic conditioned human endothelial cells. Hypertension 2001;37:1341-1348.
- 151. Nakagami H, Morishita R, Yamamoto K, Yoshimura S, Taniyama Y, Aoki M, Matsubara H, Kim S, Kaneda Y, Ogihara T. Phosphorylation of p38MAP kinase downstream of bax-caspase-3

- pathway leads to cell death induced by high D-glucose in human endothelial cells. Diabetes 2001;50:1472-1481.
- 152.Kawamura I, Morishita R (equal first author), Tsujimoto S, Manda T, Tomoi M, Tomita N, Goto T, Ogihara T, Kaneda Y. Intravenous injection of oligodeoxynucleotides to the NF-kappaB binding site inhibits hepatic metastasis of M5076 reticulosarcoma in mice. Gene Therapy 2001;8:905-912.
- 153.Sakaue G, Shimaoka M, Fukuoka T, Hiroi T, Inoue T, Hashimoto N, Sakaguchi T, Sawa Y, Morishita R, Kiyono H, Noguchi K, Mashimo T. NF-kappaB decoy suppresses cytokine expression and thermal hyperalgesia in a rat neuropathic pain model. Neuroreport. 2001;12:2079-2084.
- 154.Ahn JD, Morishita R, Kaneda Y, Lee KU, Park JY, Jeon YJ, Song HS, Lee IK, Transcription factor decoy DON for activator protein-1 (AP-1 inhibits expression of type 1 plasminogen activator inhibitor (PAI-1) gene induced by high glucose and angiotensin II in cultured human vascular smooth muscle cells. Diabetologia 2001;44:713-720.
- 155.Izumi Y, Kim S, Namba M, Yasumoto H, Miyazaki H, Hoshiga M, Kaneda Y, Morishita R, Zhan Y, Iwao H. Gene transfer of dominant-negative mutants of extracellular signal-regulated kinase and c-Jun NH2-terminal kinase prevents neointimal formation in balloon-injured rat artery. Circ Res. 2001;88:1120-1126.
- 156. Taniyama Y, Ito H, Morishita R, Ogihara T. Potential of microvascular reperfusion with adjunctive pharmacological intervention: its impact on myocardial perfusion and functional outcomes in patients with acute myocardial infarction. Drugs. 2001;61:437-441.
- 157.Aoki M, Nata T, <u>Morishita R</u>, Matsushita H, Nakagami H, Yamamoto K, Yamazaki K, Nakabayashi M, Ogihara T, Kaneda Y. Endothelial apoptosis induced by oxidative stress through activation of NFkB: anti-apoptotic effect of antioxidant agents on endothelial cells. Hypertension 2001;38:48-55
- 158.Hayashi K, Morishita R, Nakagami H, Yoshimura S, Hara A, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T, Sakai N. Gene therapy for preventing neuronal death using hepatocyte growth factor: in vivo gene transfer of HGF to subarachnoid space prevents delayed neuronal death in gerbil hippocampal CA1 neurons. Gene Therapy 2001;8:1167-1173
- 159.Otomo T. Yamamoto S, <u>Morishita R</u>, Kaneda Y. EBV replicon vector system enhances transgene expression in vivo: applications to gene therapy for cancer. The Journal of Gene Medicine. 2001;3:345-352
- 160.Sakaue G, Shimaoka M, Fukuoka T, Hiroi T, Inoue T, Hashimoto N, Sakaguchi T, Sawa Y, Morishita R, Kiyono H, Noguchi K, Mashimo T. NFkB decoy suppresses cytokine expression and thermal hyperalgesia in a rat neuropathic model. Somatosensory Systems, Pain. 2001;12:2079-2084
- 161.Aoki M, Morishita R, Hayashi SI, Jo N, Matsumoto K, Nakamaura T, Kaneda Y, Ogihara T. Inhibition of neointimal formation after balloon injury by cilostazol, accompanied by improvement of endothelial dysfunction and induction of hepatocyte growth factor in rat diabetes model. Diabetologia 2001;44:1034-1042
- 162. Taniyama Y, Morishita R, Hiraoka K, Aoki M, Nakagami H, Yamasaki K, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Therapeutic angiogenesis induced by human hepatocyte growth factor gene in rat diabetic hind limb ischemia model: molecular mechanisms of delayed angiogenesis in diabetes. Circulation 2001;104:2344-2350.
- 163. Yokoseki O, Suzuki Ji J, Kitabayashi H, Watanabe N, Wada Y, Aoki M, <u>Morishita R</u>, Kaneda Y, Ogihara T, Futamatsu H, Kobayashi Y, Isobe M. cis Element Decoy Against Nuclear Factor-kappaB Attenuates Development of Experimental Autoimmune Myocarditis in Rats. Circ Res. 2001;89:899-906.
- 164.Ueno T, Sawa Y, Kitagawa-Sakakida S, Nishimura M, <u>Morishita R</u>, Kaneda Y, Kohmura E, Yoshimine T, Matsuda H. Nuclear factor-kappaB decoy attenuates neuronal damage after global brain ischemia: A future strategy for brain protection during circulatory arrest. J Thorac

- Cardiovasc Surg. 2001;122:720-727.
- 165. Yoshimura S, <u>Morishita R</u>, Hayashi K, Yamamoto K, Nakagami H, Kaneda Y, Sakai N, Ogihara T. Inhibition of intimal hyperplasia after balloon injury in rat carotid artery model using cis-element "decoy" of nuclear factor-kB binding site as a novel molecular strategy. Gene Therapy 2001;8:1635-1642
- 166.Miyata T, Yamamoto S, Sakamoto K, <u>Morishita R</u>, Kaneda Y. Novel immunotherapy for peritoneal dissemination of murine colon cancer with macrophage inflammatory protein-1b mediated by a tumor-specific vector, HVJ catinonic liposomes. Cancer Gene Therapy 2001;8:852-860
- 167. Tsuchikane E, Nakamura T, Yamasaki K, Kobayashi T, Ptsuji S, Kobayashi T, Morishita R, Awata N. Correlation between serum hepatocyte growth factor level and percutaneous coronary balloon angioplasty: an alternative mechanisms of reduction in restenosis with cilostazol. International Journal of Angiology. 2001;10:270-274...
- 168.Namiki M, Kawashima S, Yamashita T, Ozaki M, Hirase T, Ishida T, Inoue N, Hirata Ki K, Matsukawa A, Morishita R, Kaneda Y, Yokoyama M. Local overexpression of monocyte chemoattractant protein-1 at vessel wall induces infiltration of macrophages and formation of atherosclerotic lesion: synergism with hypercholesterolemia. Arterioscler Thromb Vasc Biol. 2002;22:115-120.
- 169. Yasumoto H, Kim S, Zhan Y, Miyazaki H, Hoshiga M, Kaneda Y, <u>Morishita R</u>, Iwao H. Dominant negative c-Jun gene transfer inhibits vascular smooth muscle cell proliferation and neointimal hyperplasia in rats. Gene Therapy 2002;8:1682-1689.
- 170. Taniyama Y, Tachibana K, Hiraoka K, Namba T, Yamasaki K, Hashiya N, Aoki M, Ogihara T, Kaneda Y, Morishita R. Local delivery of plasmid DNA into rat carotid artery using ultrasound. Circulation 2002;105:1233-1239.
- 171. Morishita R, Sakaki M, Yamamoto K, Iguchi S, Aoki M, Yamasaki K, Matsumoto K, Nakamura T, Lawn R, Ogihara T, Kaneda Y. Impairment of collateral formation in Lp(a) transgenic mice: therapeutic angiogenesis induced by human hepatocyte growth factor gene. Circulation 2002;105:1491-1496
- 172. Taniyama Y, Tachibana K, Hiraoka K, Aoki M, Yamamoto S, Matsumoto K, Nakamura T, Ogihara T, Kaneda Y, <u>Morishita R</u>. Development of safe and efficient novel non-viral gene transfer using ultrasound: enhancement of transfection efficiency of naked plasmid DNA in skeletal muscle. Gene Therapy 2002;9:372-380.
- 173.Nakamura T, Morishita R, Asai T, Tsuboniwa N, Aoki M, Sakonjo H, Yamasaki K, Hashiya N, Kaneda Y, Ogihara T. Molecular strategy using cis-element "decoy" of E2F binding site inhibits neointimal formation in porcine balloon-injured coronary artery model. Gene Therapy 2002;9:488-494
- 174.Endoh M, Koibuchi N, Sato M, <u>Morishita R</u>, Kanzaki T, Murata Y, Kaneda Y. Fetal gene transfer by intrauterine injection with microbubble-enhanced ultrasound. Molecular Therapy 2002;5:501-508
- 175.Mori M, Nakagami H, <u>Morishita R</u>, Mitsuda N, Yamamoto K, Yoshimura S, Ohkubo N, Sato N, Ogihara T, Kaneda Y. N1411 mutant presenilin-2 gene enhances neuronal cell death and decreases bcl-2 expression. Life Sciences 2002;70:2567-2580.
- 176. Yoshimura S, Morishita R, Hayashi K, Kokuzawa J, Aoki M, Matsumoto K, Nakamura T, Ogihara T, Kaneda Y, Sakai N. Gene transfer of hepatocyte growth factor to subarachnoid space in cerebral hypoperfusion model. Hypertension 2002;39:1028-1034
- 177.Komai N, Ohishi M, Morishita R, Moriguchi A, Kaibe M, Matsumoto K, Rakugi H, Higaki J, Ogihara T. Serum hepatocyte growth factor concentration is correlated with the forearm vasodilator response in hypertensive patients. American Journal of Hypertension 2002;15:499-506
- 178.Ahn JD, Morishita R, Kaneda Y, Lee SJ, Kwon KY, Choi SY, Lee KU, Park JY, Moon IJ, Park JG, Yoshizumi M, Ouchi Y, Lee IK. Inhibitory Effects of Novel AP-1 Decoy

- Oligodeoxynucleotides on Vascular Smooth Muscle Cell Proliferation In Vitro and Neointimal Formation In Vivo. Circ Res. 2002;90:1325-1332.
- 179.Tomita N, Morishita R, Koike H, Hashizume M, Notake M, Fujitani B, Kaneda Y, Horiuchi M, Ogihara T. Therapeutic approach to familial hypercholesterolemia by HVJ-liposomes in LDL receptor knockout mouse. International Journal of Molecular Medicine 2002;10:137-143
- 180. Taniyama Y, Morishita R, Aoki M, Hiraoka K, Yamasaki K, Hashiya N, Matasumoto K, Nakamura T, Kaneda Y, Ogihara T. Angiogensis and anti-fibrotic action by hepatocyte growth factor in cardiomyopathic hamster. Hypertension 2002;40:47-53
- 181.Nakagami H, Morishita R, Yamamoto K, Taniyama Y, Aoki M, Yamasaki K, Matasumoto K, Nakamura T, Kaneda Y, Ogihara T. Hepatocyte growth factor (HGF) prevents endothelial cell death through inhibition of bax translocation from cytosol to mitochondrial membrane. Diabetes 2002;51:2604-2611.
- 182.Kaneda Y, Nakajima T, Nishikawa T, Yamamoto S, Ikegami H, Suzuki N, Nakamura H, Morishita R, Kotani T. Hemagglutinating Virus of Japan (HVJ) envelope vector as a versatile gene delivery system Molecular Therapy 2002;6:219-226.
- 183.Tomita N, Morishita R, Yamamoto K, Higaki J, Dzau VJ, Ogihara T, Kaneda Y. Targeted gene therapy for rat glomerulonephritis using HVJ-immunoliposomes. Journal of Gene Medicine. 2002;4:527-535
- 184.Nakamura H, Aoki M, Tamai K, Oishi M, Ogihara T, Kaneda Y, Morishita R. Prevention and regression of atopic dermatitis by ointment containing NF-kB decoy oligodeoxynucleotides in NC/Nga atopic mouse model. Gene Therapy 2002;9:1221-1229
- 185.Komai N, Morishita R, Yamada S, Ohishi M, Iguchi S, Aoki M, Sakaki M, Sakurabayashi I, Higaki J, Ogihara T. Mitogenic activity of oxidized lipoprotein (a) on human vascular smooth muscle cells. Hypertension 2002;40:310-314
- 186.Nakanishi K, Uenoyama M, Tomita N, <u>Morishita R</u>, Kaneda Y, Ogihara T, Matsumoto K, Nakamura T, Maruta A, Matsuyama S, Kawai T, Aurues T, Hayashi T, Ikeda T. Gene transfer of human hepatocyte growth factor into rat skin wounds mediated by liposomes coated with the sendai virus (hemagglutinating virus of Japan). American Journal of Pathology. 2002;161:1761-1772.
- 187.Matsumoto K, Morishita R, Tomita N, Moriguchi A, Yamasaki K, Aoki M, Matasumoto K, Nakamura T, Higaki J, Ogihara T. Impaired endothelial dysfunction in diabetes mellitus rats was restored by oral administration of prostaglandin I<sub>2</sub> analogue, accompanied by increase in vascular hepatocyte growth factor. Journal of Endocrinology 2002;175:217-223
- 188. Morishita R, Gibbons GH, Kaneda Y, Zhang L, Ogihara T, Dzau VJ. Apolipoprotein E-deficient mice created by systemic administration of antisense oligodeoxynucleotides: a new model for lipoprotein metabolism studies. Journal of Endocrinology 2002;175:475-485
- 189.Ahn JD, Morishita R, Kaneda Y, Kim HS, Chang YC, Lee KU, Park JY, Lee HW, Kim YH, Lee IK. Novel E2F decoy oligodeoxynucleotides inhibit in vitro vascular smooth muscle cell proliferation and in vivo neointimal hyperplasia. Gene Ther. 2002;9:1682-1692.
- 190. Tsukioka K, Suzuki J, Fujimori M, Wada Y, Yamaura K, Ito K, Morishita R, Kaneda Y, Isobe M, Amano J. Expression of matrix metalloproteinases in cardiac allograft vasculopathy and its attenuation by anti MMP-2 ribozyme gene transfection. Cardiovasc Res. 2002;56:472-478.
- 191.Jo N, Ogata N, Aoki M, Otsuji T, <u>Morishita R</u>, Kaneda Y, Matsumura M. Effective transfection of a cis element "decoy" of the nuclear factor-kappaB binding site into the experimental choroidal neovascularization. Curr Eye Res. 2002;24:465-473.
- 192. Shimamura M, Morishita R, Endoh M, Oshima K, Aoki M, Waguri S, Uchiyama Y, Kaneda Y. HVJ-envelope vector for gene transfer into central nervous system. Biochemical Biophysics Research Communications 2003;300:464-471
- 193.Azuma H, Tomita N, Kaneda Y, Koike H, Ogihara T, Katsuoka Y, <u>Morishita R</u>. Transfection of NFκB decoy oligodeoxynucleotides using efficient ultrasound-mediated gene transfer into donor kidneys prolonged survival of rat renal allografts. Gene Therapy 2003;10:415-425

- 194. Yamasaki K, Asai T, Shimizu M, Aoki M, Hashiya N, Sakonjo H, Makino H, Kaneda Y, Ogihara T, Morishita R. Inhibition of NFκB activation using cis-element "decoy" of NFκB binding site reduces neointimal formation in porcine balloon-injured coronary artery model. Gene Therapy 2003;10:356-364.
- 195.Koike H, Morishita R, Iguchi S, Aoki M, Matsumoto K, Nakamura T, Yokoyama C, Tanabe T, Ogihara T, Kaneda Y. Enhanced angiogenesis and improvement of neuropathy by co-transfection of human hepatocyte growth factor and prostacyclin synthase gene. The FASEB Journal 2003;17:779-781
- 196. Sekiguchi K, Yasuzumi F, <u>Morishita R</u>. Exogenous expression of hepatocyte growth factor (HGF) in rat striatum by naked plasmid DNA. Neuroscience Research 2003;45:173-180
- 197.Nakabayashi M, <u>Morishita R</u>, Nakagami H, Kuba K, Matasumoto K, Nakamura T, Kamei M, Tano Y, Kaneda Y. HGF/NK4 inhibited VEGF-induced angiogenesis in in vitro cultured endothelial cells and in vivo rabbit model. Diabetologia 2003;46:115-123
- 198.Matsumoto K, Morishita R, Tomita N, Moriguchi A, Komai N, Aoki M, Matasumoto K, Nakamura T, Higaki J, Ogihara T. Improvement of endothelial dysfunction by angiotensin II blockade, accompanied by induction of vascular hepatocyte growth factor system in diabetic spontaneously hypertensive rats. Heart & Vessels 2003;18:18-25
- 199. Yamamoto S, Yamano T, Tanaka M, Hoon D.S.B., Takao S, Morishita R, Aikou T, Kaneda Y. A novel combination of suicide gene therapy and histone deacetylase inhibitor for treatment of malignant melanoma. Cancer Gene Therapy. 2003;10:179-186.
- 200. Tomita N, Morishita R, Taniyama Y, Koike H, Aoki M, Shimizu H, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Angiogenic property of hepatocyte growth factor is dependent on up-regulation of essential transcription factor for angiogenesis, ets-1. Circulation 2003;107:1411-1417
- 201.Kato N, Nakanishi K, Nemoto K, <u>Morishita R</u>, Kaneda Y, Uenoyama M, Ikeda T, Fujikawa K. Efficient gene transfer from innervated muscle into rat peripheral and central nervous systems using a non-viral hemaggultinating virus of Japan (HVJ)-liposome method. Journal of Neurochemistry 2003;85:810-815.
- 202.Ogushi I, Iimuro Y, Seki E, Son G, Hirano T, Hada T, Tsutsui H, Nakanishi K, <u>Morishita R</u>, Kaneda Y, Fujimoto J. Nuclear factor kappa B decoy oligodeoxynucleotides prevent endotoxin-induced fatal liver failure in a murine model. Hepatology. 2003;38:335-344
- 203.Lan HY, Mu W, Tomita N, <u>Morishita R</u>, Yu XQ, Li JH, Zhu HJ, Huang XR, Johnson RJ. Inhibition of renal fibrosis by gene transfer of inducible Smad7 using ultrasound-microbubble system in rat UUO model. Journal of American Society of Nephrology 2003;14:1535-1548.
- 204.Namba T, Koike H, Murakami K, Aoki M, Makino H, Hashiya N, Ogihara T, Kaneda Y, Kohno M, Morishita R. Angiogenesis induced by endothelial nitric oxide synthase gene through VEGF expression in rat hindlimb ischemia model. Circulation 2003;108:2250-2257
- 205.Hiraoka K, Koike H, Yamamoto S, Tomita N, Yokoyama C, Tanabe T, Aikou T, Ogihara T, Kaneda Y, Morishita R. Enhanced therapeutic angiogenesis by co-transfection of prostacyclin synthase gene or optimization of intramuscular injection of naked plasmid DNA. Circulation 2003;108:2689-2696.
- 206.Kokuzawa J, Yoshimura S, Kitajima H, Shinoda J, Kaku Y, Iwama T, Morishita R, Shimazaki T, Okano H, Kunisada T, Sakai N. Hepatocyte growth factor promotes proliferation and neuronal differentiation of neural stem cells from mouse embryos. Mol Cell Neurosci. 2003;24:190-197.
- 207. Wada Y, Fujimori M, Suzuki J, Tsukioka K, Ito K, Sawa Y, <u>Morishita R</u>, Kaneda Y, Isobe M, Amano J. Egr-1 in vascular smooth muscle cell proliferation in response to allo-antigen. J Surg Res. 2003;115:294-302.
- 208.Nakashima H, Aoki M, Miyake T, Kawasaki T, Iwai M, Jo N, Oishi M, Kataoka K, Ohigi S, Ogihara T, Kaneda Y, Morishita R. Inhibition of experimental abdominal aortic aneurysm in the rat by use of decoy oligodeoxynucleotides suppressing activity of NFkB and ets

- transcription factors. Circulation 2004;109:132-138
- 209.Ohtani K, Egashira K, Usui M, Ishibashi M, Hiasa KI, Zhao Q, Aoki M, Kaneda Y, <u>Morishita R</u>, Takashita A. Inhibition of neointimal hyperplasia after balloon injury by cis-element "decoy" of early growth response gene-1 in hypercholesterolemic rabbits. Gene Therapy 2004;11:126-132.
- 210. Namiki M, Kawashima S, Yamashita T, Ozaki M, Sakoda T, Inoue N, Hirata K, <u>Morishita R</u>, Kaneda Y, Yokoyama M. Intramuscular gene transfer of interleukin-10 cDNA reduces atherosclerosis in apolipoprotein E-knockout mice. Atherosclerosis. 2004;172:21-29.
- 211. Shimamura M, Sato N, Oshima K, Aoki M, Kurinami H, Waguri S, Uchiyama Y, Ogihara T, Kaneda Y, Morishita R. A novel therapeutic strategy to treat brain ischemia: over-expression of hepatocyte growth factor gene reduced ischemic injury without cerebral edema in rat model. Circulation 2004;109:424-431
- 212.Oshima K, Shimamura M, Mizuno S, Tamai K, Doi K, Morishita R, Nakamura T, Kubo T, Kaneda Y. Intrathecal injection of HVJ-E containing HGF gene to cerebrospinal fluid can prevent and ameliorate hearing impairment in rats. FASEB J. 2004;18:212-214.
- 213.Matsuki A, Yamamoto S, Nakagami H, Aoki M, Tamai K, Matsumoto K, Nakamura T, Ogihara T, Kaneda Y, Morishita R. No influence of tumor growth by intramuscular injection of hepatocyte growth factor plasmid DNA: safety evaluation of therapeutic angiogenesis gene therapy in mice. Biochemical Biophysical Research Communications. 2004;315:59-65
- 214.Makino H, Aoki M, Hashiya N, Yamasaki K, Hiraoka K, Shimizu H, Azuma J, Kurinami H, Ogihara T, Morishita R. Increase in peripheral blood flow by intravenous administration of prostaglandin E1 in patients with peripheral arterial disease, accompanied by up-regulation of hepatocyte growth factor (HGF). Hypertension Research 2003;27:85-91
- 215. Suzuki J, Ito H, Gotoh R, Morishita R, Egashira K, Isobe M. Initial clinical cases using an NF-kB decoy at the site of the coronary stenting for prevention of restenosis. Circulation Journal 2004;68:270-271
- 216.Koibuchi N, Kaneda Y, Taniyama Y, Matsumoto K, Nakamura T, Ogihara T, <u>Morishita R</u>. Essential role of HGF (hepatocyte growth factor) in blood formation in Xenopus. Blood 2004;103:3320-3325
- 217.Hashiya N, Aoki M, Tachibana K, Taniyama Y, Yamasaki K, Hiraoka K, Makino H, Kaneda Y, Ogihara T, Morishita R. Local delivery of E2F decoy oligodeoxynucleotides using ultrasound with microbubble agent (Optison) inhibits intimal hyperplasia after balloon injury in rat carotid artery model. Biochemical Biophysical Research Communications. 2004;317:508-514.
- 218. Nakagami H, Morishita R, Nishikawa T, Shimizu H, Iguchi S, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Anti-apoptotic action of bag-1 was not related to HGF receptor, c-met, in endothelial cells. Hypertension Research 2004;27:359-365
- 219. Wada Y, Fujimori M, Suzuki J, Tsukioka K, Ito K, Sawa Y, <u>Morishita R</u>, Kaneda Y, Isobe M, Amano J. Egr-1 in vascular smooth muscle cell proliferation in response to allo-antigen. Journal of Surgical Research. 2004;115:293-302.
- 220.Hashiya N, Jo N, Aoki M, Matsumoto K, Nakamura K, Sato Y, Ogata N, Ogihara T, Kaneda Y, Morishita R. In vivo evidence of angiogenesis induced by transcription factor, ets-1: ets-1 is located upstream of angiogenesis cascade. Circulation 2004;109:3035-3041
- 221. Morishita R, Aoki M, Hashiya N, Makino H, Yamasaki K, Azuma J, Sawa Y, Matsuda H, Kaneda Y, Ogihara T. Safety evaluation of clinical gene therapy using hepatocyte growth factor to treat peripheral arterial disease by therapeutic angiogenesis. Hypertension 2004;44:203-209
- 222. YamamotoK, Tomita N, Yoshimura S, Nakagami H, Taniyama Y, Yamasaki K, Ogihara T, Morishita R. Hypoxia-induced renal epithelial cell death thourgh caspase-dependent pathway: role of bcl-2, bcl-xL and bax in tubular injury. International Journal of Molecular Medicine. 2004;14:633-640.

- 223. Yamaura K, Ito K, Tsukioka K, Wada Y, Makiuchi A, Sakaguchi M, Akashima T, Fujimoto M, Sawa Y, Morishita R, Matsumoto K, Nakamura T, Suzuki J, Amano J, Isobe M. Suppression of acute and chronic rejection by hepatocyte growth factor in a murine model of cardiac transplantation: induction of tolerance and prevention of cardiac allograft vasculopathy. Circulation 2004;110:1650-1657.
- 224.Katsuragi N, <u>Morishita R</u>, Nakamura N, Ochiai T, Taniayama Y, Hasegawa Y, Kawashima K, Kaneda Y, Ogihara T, Sugimura K. Periostin as a novel factor responsible for ventricular dilation. Circulation 2004;110:1806-1813
- 225. Shimamura M, Sato N, Taniyama Y, Yamamoto S, Endoh M, Kurinami H, Aoki M, Ogihara T, Kaneda Y, Morishita R. Development of efficient plasmid DNA transfer into adult rat central nervous system using microbubble-enhanced ultrasound. Gene Therapy 2004;11:1532-1539
- 226.Hiraoka K, Yamamoto S, Otsuru S, Nakai S, Tamai K, Morishita R, Ogihara T, Kaneda Y. Enhanced tumor-specific long-term immunity of Hemagglutinating Virus of Japan-mediated dendritic cell-tumor fused cell vaccination by coadministration with CpG oligodeoxynucleotides. Journal of Immunology 2004;173:4297-4307.
- 227. Yamasaki K, Aoki M, Makino H, Hashiya N, Shimizu H, Ohishi M, Ogihara T, <u>Morishita R</u>. Effect of nifedipine on endothelial function in normotensive smokers: potential contribution of increase in circulating hepatocyte growth factor. Journal of Human Hypertension 2004;28:701-705
- 228.Powell RJ, Dormandy J, Simons M, <u>Morishita R</u>, Annex BH. Therapeutic angiogenesis for critical LIMB ischemia: design of the haptocyte growth factor therapeutic angiogenesis clinical trial. Vascular medicine 2004;9:1-6
- 229. Tomita N, Kim JY, Gibbons GH, Zhang L, Kaneda Y, Stahl RA, Ogborn M, Venderville B, Morishita R, Baran D, Dzau VJ. Gene therapy with an E2F transcription factor decoy inhibits cell cycle progression in rat anti-Thy 1 glomerulonephritis. Int J Mol Med. 2004;13:629-636
- 230.Ahn JD, Morishita R, Kaneda Y, Kim HJ, Kim YD, Lee HJ, Lee KU, Park JY, Kim YH, Park KK, Chang YC, Yoon KH, Kwon HS, Park KG, Lee IK. Transcription factor decoy for AP-1 reduces mesangial cell proliferation and extracellular matrix production in vitro and in vivo. Gene Ther. 2004;11:916-923.
- 231.Koike H, Tomita N, Azuma H, Taniyama Y, Yamasaki K, Kunugiza Y, Tachibana K, Ogihara T, **Morishita R**. An efficient gene transfer method mediated by ultrasound and microbubbles into the kidney. Journal of Gene Medicine 2005;7:108-116.
- 232.Kato N, Nemoto K, Nakanishi K, <u>Morishita R</u>, Kaneda Y, Uenoyama M, Ikeda T, Fujikawa K. Nonviral gene transfer of human hepatocyte growth factor improves streptozotocin-induced diabetic neuropathy in rats. Diabetes. 2005;54:846-854.
- 233. Miwa K, Nakashima H, Aoki M, Miyake T, Kawasaki T, Iwai M, Oishi M, Kataoka K, Ohigi S, Ogihara T, Kaneda Y, <u>Morishita R</u>. Inhibition of ets, an essential transcription factor for angiogenesis, to prevent the development of abdominal aortic aneurysm in a rat model. Gene Therapy 2005;12:1109-1118.
- 234.Makino H, Aoki M, Hashiya N, Yamasaki K, Shimizu H, Ogihara T, <u>Morishita R</u>. A calcium channel blocker, benidipine, improves forearm reactive hyperemia in patients with essential hypertension. Blood Pressure Suppl. 2005;1:39-44.
- 235. Yumoto A, Fukushima Kusano K, Nakamura K, Hashimoto K, Aoki M, <u>Morishita R</u>, Kaneda Y, Ohe T. Hepatocyte growth factor gene therapy reduces ventricular arrhythmia in animal models of myocardial ischemia. Acta Med Okayama. 2005;59:73-78.
- 236.Kato N, Nemoto K, Nakanishi K, <u>Morishita R</u>, Kaneda Y, Uenoyama M, Ikeda T, Fujikawa K. Nonviral HVJ (hemagglutinating virus of Japan) liposome-mediated retrograde gene transfer of human hepatocyte growth factor into rat nervous system promotes functional and histological recovery of the crushed nerve. Neurosci Res. 2005;52:299-310.
- 237. Tashiro H, Aoki M, Isobe M, Hashiya N, Makino H, Kaneda Y, Ogihara T, <u>Morishita R</u>. Development of novel method of non-viral efficient fene transfer into neonatal cardiac

- myocytes. Journal of Molecular and Cellular Cardiology 2005;39:503-509
- 238.Morishita N, Nakagami H, Morishita R, Takeda S, Mishima F, Tarazono B, Nishijima S, Kaneda Y, Tanaka N. Magnetic nanoparticles with surface modification enhanced gene delivery of HVJ-E vector. Biochemical Biophysical Research Communications 2005;334:1121-1126.
- 239. Taniyama Y, Kurinami H, Tanaka H, Takami T, Ogihara T, Tohyama M, Kaneda Y, Morishita R. Gene transfer into adult rat spinal cord using naked plasmid DNA and ultrasound microbubbles. Journal of Gene Medicine 2005;7:1468-1474
- 240.Nakagami H, Maeda K, <u>Morishita R</u>, Iguchi S, Nishikawa T, Takami Y, Kikuchi Y, Saito Y,. Tamai K, Ogihara T, Kaneda Y. Novel autologous cell therapy in ischemic limb disease through growth factor secretion by cultured adipose tissue-derived stromal cells. Artrioscleros, Thrombosis and Vascular Biology 2005;25:2542-2547.
- 241 Inagaki H, Suzuki J, Ogawa M, Taniyama Y, <u>Morishita R</u>, Isobe M. Ultrasound-microbubble mediated NF-kB decoy transfection attenuates neointimal formation after arterial injury in mice. Journal of Vascular Research 2006;43:12-18
- 242.Kim YD, Park KG, Morishita R, Kaneda Y, Kim SY, Song DK, Kim HS, Nam CW, Lee HC, Lee KU, Park JY, Kim BW, Kim JG, Lee IK. Liver-directed gene therapy of diabetic rats using an HVJ-E vector containing EBV plasmids expressing insulin and GLUT 2 transporter. Gene Therapy 2006;13:216-224.
- 243.Miyake T, Aoki M, Shiraya S, Kawasaki T, Oishi M, Kataoka K, Tanemoto K, Ogihara T, Kaneda Y, Morishita R. Prevention of aneurysms by simultaneous inhibition of NFkB and ets using chimeric decoy oligonucleotides in rabbit model. Gene Therapy 2006;13:695-704
- 244. Shimamura M, Sato N, Waguri S, Uchiyama Y, Hayashi T, Iida H, Nakamura T, Ogihara T, Kaneda Y, Morishita R. Gene transfer of HGF gene improves learning and memory in the chronic stage of cerebral infarction. Hypertension 2006;47:742-751
- 245. Koibuchi N, Taniyama Y, Nagao K, Ogihara T, Kaneda Y, <u>Morishita R</u>. The Effect of VEGF on blood vessels and blood cells during <u>Xenopus</u> development. Biochemical Biophysical Research Communications 2006;344:339-345
- 246.Shimizu H, Nakagami H, Tsukamoto I, Morita S, Kunugiza Y, Tomita T, Yoshikawa H, Kaneda Y, Ogihara T, Morishita R. NFkB decoy oligodeoxynucleotides ameliorates osteoporosis through inhibition of activation and differentiation of osteoclasts. Gene Therapy 2006;13:933-941
- 247.Nishikawa T, Nakagami H, Matsuki A, Maeda A, Yo CY, Harada T, Morishita R, Tamai K, Kaneda Y. Development of high throughput functional screening of therapeutic genes using HVJ-E vector. Human Gene Therapy 2006;17:470-475
- 248.Sato N, Okochi M, Taniyama Y, Kurinami H, Shimamura H, Takeuchi D, Hamada H, Fukumori A, Kiyosue K, Taguchi T, Tanaka T, Miyasaki M, Takeda M, Ogihara T, Morishita R. Development of nNew sScreening sSystem for Alzheimer Disease, iIn vVitro Aβ sSink aAssay, to iIdentify the dDissociation of soluble Aβ 42 is more resistant to dissociate dissociation ffFrom fibrilsfFibrils. Neurobiology of Disease 2006;22:487-495.
- 249.Nakagami H, Nakagawa N, Takeya Y, Kashiwagi K, Ishida C, Hayashi S, Aoki M, Matsumoto K, Nakamura T, Ogihara T, <u>Morishita R</u>. A model of vasculogenesis from ES cells for vascular research and regenerative medicine. Hypertension 2006;48:112-119
- 250.Kunugiza Y, Tomita N, Taniyama Y, Tomita T, Mariana KO, Tamai K, Tanabe T, Kaneda Y, Yoshikawa H, Morishita R. Acceleration of wound healing by combined gene transfer of hepatocyte growth factor and prostacycline synthase with shima jet. Gene Therapy 2006;13:1143-1152
- 251.Tomita T, Kunugiza Y, Tomita N, Takano H, <u>Morishita R</u>, Kaneda Y, Yoshikawa H. E2F decoy oligodeoxynucleotide ameliorates cartilage invasion by infiltrating synovium derived from rheumatoid arthritis. International Journal of Molecular Medicine 2006;18:257-265
- 252. Azuma H, Taniyama Y, Takeya Y, Iekushi K, Aoki M, Dosaka N, Matsumoto K, Nakamura T,

- Ogihara T, Morishita R. Angiogenic and anti-fibrotic actions of hepatocyte growth factor improve cardiac dysfunction in porcine ischemic cardiomyopathy. Gene Therapy 2006;13:1206-1213.
- 253.Saito Y, Nakagami H, Morishita R, Takami Y, Kikuchi Y, Hayashi H, Nishikawa T, Tamai K, Azuma N, Sasajima T, Kaneda Y. Transfection of human hepatocyte growth factor gene ameliorates secondary lymphedema via promotion of lymphangiogenesis. Circulation 2006;114:1177-1184
- 254. Miyake T, Aoki M, Shiraya S, Tanemoto K, Ogihara T, Kaneda Y, Morishita R. Inhibitory effects of NFkB decoy oligodeoxynucleotides on neointimal hyperplasia in a rabbit vein graft model. Journal of Molecular and Cellular Cardiology (in press)
- 255.Koike H, Ishida A, Shimamura M, Mizuno S, Nakamura T, Ogihara T, Kaneda Y, <u>Morishita</u> R. Prevention of onset of Parkinson's disease by <u>in vivo</u> gene transfer of human hepatocyte growth factor in rodent model: a model of gene therapy for Parkinson's disease. Gene Therapy (in press)
- 256. Shiraya S, Miwa K, Aoki M, Miyake T, Oishi M, Kataoka K, Ohgi S, Ogihara T, Kaneda Y, Morishita R. Hypertension accelerated experimental abdominal aortic aneurysm through up-regulation of transcription factors, NFκB and Ets. Hypertension (in press)
- 257. Kunugiza Y, Tomita T, Tomita N, <u>Morishita R</u>, Yoshikawa H. Inhibitory effect of ribbon-type NF-kappaB decoy oligodeoxynucleotides on osteoclast induction and activity in vitro and in vivo. Arthritis Res Ther. 2006;8:R103 (in press)

#### Reviews

- 1. Morishita R, Gibbons GH, Dzau VJ. Analysis of cellular uptake of antisense oligonucleotides using FITC labeled oligonucleotides. Oligotechniques 1994;2:1-3.
- 2. Dzau VJ, Gibbons GH, <u>Morishita R</u>, Pratt E. New perspectives in hypertension research: potentials of vascular biology. Hypertension 1994;23:1132-1140.
- 3. Dzau VJ, <u>Morishita R</u>, Gibbons GH. Cardiovascular gene therapy: possibilities and realities. Proceedings of the 3rd International Symposium On Multiple Risk Factors In Cardiovascular Disease Vascular And Organ Protection (in press)
- 4. <u>Morishita R</u>, Gibbons GH, Dzau VJ. Potential for transcatheter application of antisense oligonucleotides for treatment of vascular diseases. Journal of Interventinal Cardiology 1995;8:377-381.
- 5. Kaneda Y, and Morishita R. In vivo gene transfer for the therapy of vascular diseases. Journal of Atherosclerosis Society. 1995;22:819-823.
- 6. Kaneda Y, <u>Morishita R</u>, Tomita N. Increased expression of DNA cointroduced with nuclear protein in adult rat liver. Journal of Molecular Medicine 1995:73:289-297.
- 7. Morishita R, Higaki J, Aoki M, Hayashi S, Kida I, Kaneda Y, Ogihara T. Novel strategy of gene therapy in cardiovascular disease with HVJ-liposome method. In "Progression of chronic renal diseases", Koide H (eds.) Contribution to Nephrology, 1996;118:254-264.
- 8. Dzau VJ, Mann MJ, <u>Morishita R</u>, Kaneda Y. Fusigenic viral liposome for gene therapy in cardiovascular diseases. Proc Natl Acad Sci USA 1996;93:11421-11425.
- 9. Morishita R, Aoki M, Nakamura S, Matsushita H, Tomita N, Hayashi S, Moriguchi A, Matsumoto K, Nakamura T, Ogihara T. Potential role of a novel vascular modulator, hepatocyte growth factor (HGF), in cardiovascular disease; characterization and regulation of local HGF system. Journal of Atherosclerosis and Thrombosis 1997;4:12-19.
- 10. Tomita N, <u>Morishita R</u>, Higaki J, Ogihara T. Strategy for functional inactivation of genes:a novel strategy for gene therapy and gene regulation analysis using transcriptional factor decoy oligonucleotides. Experimental Nephrology 1997;5:429-434.
- 11. Kaneda Y, Morishita R, Dzau VJ. Prevention of restenosis by gene therapy. Ann NY Acad Sci 1997;811:299-308
- 12. Mann MJ, Morishita R, Gibbons GH, von-der-Leyen HE, Dzau VJ. DNA transfer into

- vascular smooth muscle using fusigenic Sendai virus (HVJ)-liposomes. Molecular Cellular Biochemstry 1997;172:3-12.
- 13. <u>Morishita R</u>, Ogihara T. <u>In vivo</u> evaluation of <u>in vitro</u> hypothesis using "gain or loss" of function: functional analysis of renin-angiotensin system. American Journal of Hypertension 1998;11:507-513.
- 14. Morishita R, Nakamura S, Hayashi S, Aoki M, Matsushita H, Tomita N, Yamamoto K, Moriguchi A, Higaki J, Ogihara T. Contribution of a vascular modulator, hepatocyte growth factor (HGF), to the pathogenesis of cardiovascular disease. Journal of Atherosclerosis and Thrombosis 1998;4:129-134..
- 15. <u>Morishita R</u>, Higaki J, Tomita N, Ogihara T. Application of transcription factor "decoy" strategy as means of gene therapy and study of gene expression in cardiovascular disease. Circulation Research 1998;82:1023-1028
- 16. <u>Morishita R</u>. Lessons from human arteries: how to design a gene therapy strategy for treatment of cardiovascular disease. (Editorial) Circulation Research 1998;82:1349-1351
- 17. <u>Morishita R</u>, Nakagami H, Taniyama Y, Matsushita H, Yamamoto K, Tomita N, Moriguchi A, Matsumoto K, Higaki J, Ogihara T. Oligonucleotide-based gene therapy for cardiovascular disease. Clinical Chemistry and Laboratory Medicine 1998;36:529-534
- 18. Dzau VJ, von-der-Leyen HE, <u>Morishita R</u>. The concept and potentials of cardiovascular gene therapy. Indian Heart J. 1998;50:23-33
- 19. Kaneda Y, Saeki Y, Morishita R. Gene therapy using HVJ-liposomes: the best of both worlds? Molecular Medicine Today 1999;7:298-303.
- 20. <u>Morishita R</u>, Moriguchi A, Higaki J, Ogihara T. Hepatocyte growth factor as potential index of severity of hypertension. Hypertension Research 1999;22:161-167
- 21. <u>Morishita R</u>, Aoki M, Ogihara T. Functional analysis of cardiovascular renin-angiotensin system using "gain or loss" of function approach. Hypertension Research 2000;23:137-141
- 22. <u>Morishita R</u>, Aoki M, Kaneda Y. Oligonucleotides-based gene therapy for cardiovascular disease: are oligonucleotides therapeutics novel cardiovascular drugs? Current Drug Targets 2000;1:15-23.
- 23. Tomita N, <u>Morishita R</u>, Kaneda Y, Higaki J, Ogihara T. Gene therapy as a potential treatment for restenosis and myocardial infarction. Drug News & Perspectives 2000;13:206-212.
- 24. Aoki M, <u>Morishita R</u>, Taniyama Y, Kaneda Y, Ogihara T. Therapeutic angiogenesis induced by hepatocyte growth factor: potential gene therapy for ischemic diseases. J Atheroscler Thromb. 2000;7:71-76.
- 25. Tomita N, Morishita R, Higaki J, Ogihara T. Novel molecular therapeutic approach to cardiovascular disease based on hepatocyte growth factor. J Atheroscler Thromb. 2000;7:1-7.
- 26. Morishita R, Aoki M, Kaneda Y, Ogihara T. Gene Therapy in vascular medicine: recent ad vances and future perspectives. Pharmacology & Therapeutics 2001;91:105-114
- 27. Nakamura H, Morishita R, Kaneda Y. Molecular therapy via transcriptional regulation with double-stranded oligodeoxynucleotides as decoys. In Vivo 2002;16:45-48
- 28. Tomita N, Morishita R, Tomita T, Ogihara T. Potential therapeutic applications of decoy oligonucleotides. Current Opinion in Molecular Therapeutics 2002;4:166-170
- 29. Morishita R, Kaneda Y. HVJ (hemagglutinating virus of Japan; Sendai virus)-liposome method. Methods Enzymol. 2002;346:619-27.
- 30. <u>Morishita R</u>, Aoki M, Yo Y, Ogihara T. Hepatocyte growth factor as "cardiovascular hormone": role of HGF in the pathogenesis of cardiovascular disease. Endocrine Journal 2002;49:273-284
- 31. Morishita R, Aoki M, Hashiya N, Yamasaki K, Makino H, Wakayama K, Azuma J, Ogihara T. Hepatocyte growth factor (HGF) angiogenic gene therapy: promises for cardiovascular diseases Gene Therapy & Regulation 2002;1:343-359
- 32. Morishita R, Tomita N, Ogihara T. HMG-Co A reductase inhibitors in the treatment of cardiovascular diseases: stabilization of coronary artery plaque. Current Drug Targets

- 2002;3:379-385
- 33. Morishita R, Ogihara T. Gene therapy for peripheral arterial occlusive disease. Japan Medical Association Journal 2002;45:496-498
- 34. Morishita R. Recent progress in gene therapy to treat cardiovascular disease. Circulation Journal 2002;66:1077-1086
- 35. Tomita N, Morishita R, Tomita T, Azuma H, Ogihara T. NF-kB as a target for transcription factor decoy strategy in inflammatory diseases. Current Medicinal Chemistry 2003;2:51-56
- 36. Tomita N, Azuma H, Kaneda Y, Ogihara T, <u>Morishita R</u>. Gene therapy with transcription factor decoy oligonucleotides as a potential treatment for cardiovascular diseases. Current Drug Targets 2003;4:339-346
- 37. Morishita R. Cardiovascular disease and angiogenesis. Internal Medicine. 2003;42:301-302
- 38. Tomita N, Tomita T, Yuyama K, Tougan T, T Tajima, Ogihara T, Morishita R. development of novel decoy oligonucleotides: adavantages of circular dumb-bell decoy. Current Opinion in Molecular Therapeutics. 2003;5:107-112
- 39. Morishita R, Aoki M, Kaneda Y, Ogihara T. Therapeutic potential of oligonucleotides-based therapy in cardiovascular disease. BioDrugs 2003;17:383-389.
- 40. <u>Morishita R</u>. Transcription factor as molecular targets: is transcription factor decoy a novel drug? Current Drug Target 2003;4:2 p before 599
- 41. Tomita N, Morishita R, Ogihara T. Transcription factors as molecular targets: molecular mechanisms of decoy ODN and their design. Current Drug Target 2003;4:603-608
- 42. Aoki M, Morishita R, Ogihara T. Oligonucleotide-based molecular therapy for restenosis sfter angioplasty. Current Drug Target 2003;4:613-618.
- 43. Morishita R, Tomita N, Kaneda Y, Ogihara T. Molecular therapy to inhibit NFkB activation by transcription factor decoy oligonucleotides. Current Opinion in Pharmacology 2004;4:139-146
- 44. Kotani H, Nakajima T, <u>Morishita R</u>, Kaneda Y. Innovative novel vectors for cardiovascular disease. Current Gene Therapy 2004;4:183-194
- 45. Morishita R, Aoki M, Hashiya N, Yamasaki K, Kurinami H, Shimizu S, Makino H, Takeya Y, Azuma J, Ogihara T. Therapeutic angiogenesis using hepatocyte growth factor (HGF). Current Gene Therapy 2004;4:199-206
- 46. Morishita R. Perspective in progress of cardiovascular gene therapy. Journal of Pharmacological Science 2004;95:1-8
- 47. <u>Morishita R</u>. Is VEGF a missing link between hypertension and inflammation? Hypertension (Editorial) 2004;44:253-254
- 48. Nakagami H, Kaneda Y, Ogihara T Morishita R. Endothelial dysfunction in hyperglycemia as a trigger of atherosclerosis. Current Diabetes Review 2005;1:59-63
- 49. Nakagami H, Kaneda Y, Ogihara T, <u>Morishita R</u>. Hepatocyte growth factor as potential cardiovascular therapy. Expert Review of Cardiovascular Therapy 2005;3:513-519
- 50. Morishita R, Aoki M, Ogihara T. Does gene therapy become pharmacotherapy? Experimental Physiology 2005;3:307-313
- 51. Sato N, Shimamura M, <u>Morishita R.</u> Recent progress in cerebrovascular gene therapy. Current Neurovascular Research 2005;2:235-247
- 52. Morishita R. Scientific rationale of the CREST trial results: evidence for the mechanism of action of cilostazol in restenosis. Atherosclerosis 2005;6:41-46.
- 53. Shimamura M, Sato N, Yoshimura S, Kaneda Y, <u>Morishita R.</u> HVJ-based non-viral gene transfer method: successful gene therapy using HGF and VEGF genes in experimental ischemia. Frontiers in Bioscience 2006;11:753-759
- 54. Nakagami H, Tomita N, Kaneda Y, Ogihara T, <u>Morishita R.</u>. Endothelial dysfunction in hyperglycemia as a trigger of atherosclerosis: anti-oxidant gene therapy by NFkB decoy oligonucleotide. Current Pharmaceutical Biotechnology 2006;7:95-100.
- 55. Nakagami H, Morishita R, Maeda K, Kikuchi Y, Ogihara T, Kaneda Y.Adipose tissue-derived

- stromal cells as a novel option for regenerative cell therapy. Journal of Atherosclerosis and Thrombosis. 2006;13:77-81.
- 56. Morishita R, Tomita N, Ogihara T. Statin biology beyond cholesterol lowering: vascular statin vs. hepatic statin. Expert Review of Cardiovascular Therapy (in press)
- 57. <u>Morishita R</u>. Gene Therapy in cardiovascular medicine: recent advances and perspectives. Drugs of Today (in press)
- 58. Tomita N, <u>Morishita R</u>. Antisense oligonucleotides as a powerful molecular strategy for gene therapy in cardiovascular diseases Current Pharmaceutical Design (in press)
- 59. Kawakami K, Nakajima O, <u>Morishita R</u>, Nagai R. Targeted anticancer immunotoxins and cytotoxic agents with direct killing moieties. The Scientific World Journal 2006;6:781-90.

#### Book

- 1. Higaki J, Ogihara T, Nakamaru M, Mikami M, Morishita R, Kumahara Y. Hypothalamic-pituitary-adrenal axis in primary aldosteronism. In:The adrenal and hypertension: from cloning to clinic. Serono symposium publications 57,201-207, eds Mantero S, Takeda R, Scoggins BA, Opocher G, Raven Press, NY.
- 2. <u>Morishita R</u>, Gibbons GH, Dzau VJ. Gene therapy as potential treatment for cardiovascular diseases. Cardiovascular Pharmacology And Therapeutics eds, Singh BN, p51-61, New Tork, Libingstone Publisher, 1993
- 3. Yamada K, Moriguchi A, Mikami H, Higaki J, Morishita R, Aoki M, Okuda N, Ogihara T. Importance of central amino acid neurotransmitter in the hypotensive effect of angiotensin II type I receptor antagonist in the spontaneously hypertensive rat. Central Nervous System and Blood Pressure Control, edited by Niwa M, Imaizumi T and Takahashi H, Yubunsha, Tokyo, Japan, 1995, 65-70.
- 4. Dzau VJ, Von Der Leyen HE, <u>Morishita R</u>. The concept and potentials of cardiovascular gene therapy. In: Dialogues in Cardiovascular Medicine; Gene Therapy 1997;2.1-3.
- 5. Morishita R, Aoki M, Matsumoto H, Hayashi S, Nakamura S, Nakano N, Nishii T, Yamamoto K, Tomita N, Moriguchi A, Ogihara T. Functional analysis of tissue renin-angiotensin System using "gain and loss of function" approaches: in vivo test of in vitro hypothesis. in "Angiotensin II receptor blockade: physiological and clinical implications" edited by Dhalla NS, Zahradka P, Dixon IMC, Beamish RE. Kluwer Academic Publishers, Boston, 1997. (in press)
- 6. Morishita R, Aoki M, Matsushita H, Kaneda Y, Higaki J, Ogihara T. Gene therapy for myocardial infarction. The Ischemic Heart edited by Mochizuki S, Takeda N, Nagano M, Dhalla NS. Kluwer Academic Publishers, Boston, 1998. 531-550.
- 7. Morishita R, Aoki M, Nakamura S, Nakagami H, Taniyama Y, Yamamoto K, Moriguchi A, Higaki J, Ogihara T, Kaneda Y. Gene therapy for cardiovascular disease: is gene therapy a real medicine in next millennium? Proceeding of 3rd Green Cross International Symposium, Edited by Suzuki K, Ikeda Y, Maruyama I. Eibun Press Ltd., Osaka, 2000:217-229.
- 8. Morishita R, Aoki M, Nakagami H, Taniyama Y, Yamamoto K, Matsushita H, Matsumoto K, Higaki J, Kaneda Y, Ogihara T. Gene therapy for myocardial infarction: a real medicine in next millenium? Heart Failure; Frontiers in Cardiology, eds. Kitabatake A, Sasayama S, Francis G.S. 2000, Springer, Tokyo, 219-226
- 9. Isobe M, Suzuki J, <u>Morishita R</u>, Kaneda Y, Amano J. Gene therapy for heart transplantation-associated coronary arteriosclerosis. Atherosclerosis V. The Fifth Saratoga International Conference. Edited by Numano F & Gimbrone MA. Ann New York Acad Sci 2000, 902: 77-83.
- 10. <u>Morishita R</u>, Aoki M, Nakamura S, Higaki J, Kaneda Y, Ogihara T. Gene therapy for cardiovascular disease using hepatocyte growth factor. Atherosclerosis V. The Fifth Saratoga International Conference. Edited by Numano F & Gimbrone MA. Ann New York Acad Sci

- 2000, 902: 369-376.
- 11. Kaneda Y, Nakasima H, Tsuboniwa N, <u>Morishita R</u>. Progress of fusigenic viral-liposomes. Progress in Gene Therapy: basic and clinical frontiers. edited by Bertolottis R et al. VSP, 2000:225-243.
- 12. Morishita R, Aoki M, Kaneda Y. Decoy oligodeoxynucleotides as novel cardiovascular drugs for cardiovascular disease. Atherosclerosis VI. Edited by Bertolottis R et al. Numano F and MA Gimbrone. Annals of The New York Academy of Sciences. 2001;947:294-302
- 13. Morishita R, Kaneda Y. HVJ-liposome method. "Methods in Enzymology" edited by M I Phillips. Academic Press. 2002, 619-627
- 14. <u>Morishita R</u>. Cardiovascular disease: gene therapy. The Encyclopedia of the Human Genome, Nature Publishing Group. 2003(in press)
- 15. Morishita R, Tomita N, Kaneda Y, Ogihara T. Potential of transcription factor decoy oligonucleotides as therapeutic approach. "Transcription Factor" in "Handbook of Experimental Pharmacology", edited by M. Gossen, S. Triezenberg and J. Kaufmann, the Springer-Verlag. 2004; 439-453
- 16. Morishita R, Tomita N, Ogihara T. Decoy oligodeoxynucleotides as viable pharmaceutical drugs. "Synthetic nucleic acids as inhibitors of gene expression: mechanisms, application and therapeutic implications" edited by Dr. Levon Khachigian, Nova Science (Hauppauge, NY) 2004:161-175
- 17. Morishita R. Gene therapy vs pharmacotherapy. "Contemporary Cardiology: Cardiovascular genomics" edited by Dr. M/K/ Raizada et al. Humana Press Inc., Totowa, NJ. 2005:137-156

18. Symposium:


1. Featured Research symposia "Therapy of restenosis - Molecular straegies", 65th Scientific Sessions of American Heart Association, New Orleans, Nov 17,1992.


"In vivo gene transfer into intact blood vessels: a bovel and efficienct method" Morishita R, Gibbons GH, Zhang L, Kaneda Y, Ogihara T, Dzau VJ.

2. Featured Research symposia "Atherosclerosis and Vascular biology", American Fedaration of Clinical Research, Washington, May 1, 1993.

"Novel intraluminal molecular delivery of antisense cdc 2 kinase and PCNA oligonucleotides inhibits neointimal hyperplasia"

Morishita R, Gibbons GH, Ellison KE, Nakajima M, Zhang L, Kaneda Y, Ogihara T, Dzau VJ.





EU-RUM IN 1116

# STEDMAN'S

# Medical Dictionary

28th Edition

Illustrated in Color



LIPPINCOTT WILLIAMS & WILKINS

A Wolters Kluwer Company

Philadelphia • Baltimore • New York • London Buenos Aires • Hong Kong • Sydney • Tokyo Publisher: Julic K. Stegman
Senior Product Manager: Eric Branger
Associate Managing Editor: Tiffany Piper
Marketing Manager: Yvonne Palmer
New Terms Editor: Thomas W. Filardo, MD

New Terms Euror. Inch. Direkx, MD
Etymology Editor: John H. Direkx, MD
Chief Copyeditor: Raymond Lukens

Copyeditors: Ellen Atwood, Vincent Ercolano, Ellen Erkess, Linda Scaring

Pronunciations Editor: William R. Hensyl

On-Line Editors: Barbara L. Ferretti, Kathryn J. Cadle, Lisa Fahnostock

Proofreaders: Richard Diamanti, Raymond Lukons, Kristi Lukons, Joff Prucher, Linda C. Robbian

Database Programmers: Dave Marcus, Lexi-Comp Inc., Hudson, OH

Art Director: Jennifer Clements Illustrations: Neil O. Hardy

Additional artwork by: Mary Anna Barratt Dimes, Kathryn Born, Rob Duckwall, Timothy Hengst, Mikki Senkarik, Michael Schenk, Latry Ward

Graphic Artist: Susan Caldwell
Typographic Design: Parkton Art Studio, Inc.

Copyright © 2006
Lippincott Williams & Wilkins
A Wolters Kluwer Health Company
351 West Camden Street
Builtimore, Maryland 21201-2436 USA
www.stedmans.com
stedmans@iww.com

Copyright © by William Wood and Company: 1911, 1st ed.; 1912, 2nd ed.; 1914, 34 ed.; 1916, 4th ed.; 1918, 5th ed.; 1920, 6th ed.; 1922, 7th ed.; 1924, 8th ed.; 1926, 9th ed.; 1928, 10th ed.; 1930, 11th ed.

Copyright © by Williams & Wildens: 1933, 12th ed; 1935, 13th ed.; 1939, 14th ed.; 1942, 15th ed.; 1946, 16th ed.; 1949, 17th ed.; 1953, 18th ed.; 1957, 19th ed.; 1961, 20th ed.; 1966, 21st ed.; 1972, 22nd ed.; 1976, 23rd ed.; 1982, 24th ed.; 1990, 25th ed.; 1995, 26th ed.

Copyright © by Lippincott Williams & Wilkins: 2000, 27th cd.

All rights reserved. This book and software are protected by copyright. No part of this book or software may be reproduced in any form or by any means, including photocopying, or utilized by any information storage and retrieval system without written permission from the copyright owner.

Stedman's, STEDMAN'S is a registered trademark of Lippincott Williams & Wilkins.

The publisher is not responsible (as a matter of product liability, negligence or otherwise) for any injury resulting from any material contained hetein. This publication contains information relating to general principles of medical care which should not be construed as specific instructions for individual patients. Manufacturers' product information and package inserts should be reviewed for current information, including contraindications, desages and premutions.

Datubase design by Lexi-Comp Inc. Hudson, OH Printed in the United States of America by Quebecor World

Library of Congress Cataloging-in-Publication Data Stedman, Thomas Lathrop, 1853-1938,

Stedman's medical dictionary. -28th ed.

p.; cm.
Includes bibliographical references and index.
ISBN 0-7817-3390-1 (alk, paper)
1. Medicine—Dictionatics, I. Title, II. Title: Medicul dictionary.
[DNLM: 1. Medicine—Dictionary—English. W 13 S812m 2006]
R121-S8 2006
610'3—dc22

2005021544

#### bradycardiac

250

浅村内外特許事務所

branci

brad-y-car-di-ac (brad'e-km'de-ak). Relating to or characterized by bradycardin syn bradycardic.

bra.dy.car.dlc (brad's-kar'dlk). syn bradycardiae.

bra-dy-cl-ne-sl-a (brad'ē-si-ne'sē-š). syn bradykincsia.

bra-dy-crot-ic (brad'ë-krot'ik). Relating to or characterized by a slow pulse. [brady + G. krotos, a striking]

bra-dy-di-as-to-le (brad'ē-dī-as'tō-lē). Prolongation of the disstole of the heart.

bra-dy-es-the-si-a (brad'ē-es-thē'zē-š). Slow sensory per-ception. [brady-+ G. aisthēsis, sensation]

bradygastria (brad-ē-gas'trē-ā). Decreased rate of electrical pacemaker activity in the stomach, defined as less than a cycles/minute for at least I minute. Normal activity is defined as an electrical signal at a frequency of 2-4 cycles per minute using cutaneous clectrogastrography. May be associated with nausea, gastroparesis, irritable bowel syndrome, and functional dyspepsia,

bra-dy-glos-si-a (brad'ē-glos'ē-ā). 1. Slow or difficult tongue movement 2. syn bradyarthria. [brady- + G. glossa, tongue]

bra-dy-ki-ne-si-a (brud'ē-kin-ē'zē-ii). A decrease in spontancity and movement. One of the features of extrapyramidal disorders, such as Parkinson disease, syn bradycinesia. [brady-+ G. kinēsis, movement)

bra-dy-ki-net-ic (brad'c-ki-net'ik). Characterized by or pertaining to slow movement.

bra-dy-kl-nin (brad'e-kt'nin). The nonapeptide Arg-Pro-Pro-Gly-Phe-Ser-Pro-Phe-Arg, produced from the decapentide kalli-din (bradykininogen) that is produced from oz-globulin by kalli-krein, normally present in blood in an inactive form and similar to trypsin in action; b. is one of several plasma kinins, is a potent vasodilator, and is one of the physiologic mediators of anaphylaxis released from cytotropic antibody-coated mast cells following reaction with antigen (allergen) specific for the antibody. sankallidin 9, kallidin 1, kinin 9. [brady- + G, kineū, to move]

bra-dy-ki-nin-o-gen (brad'ē-ki-nin'ō-jen). Nyn kallidin.

bra-dy-ki-nin po-ten-ti-a-tor B (brad'e-ki nin po-ten'sheä-tör). Glp-Gly-Leu-Pro-Pro-Arg-Pro-Lys-Ile-Pro-Pro; the undecapeptide precursor of bradykinin and the angiotensins,

bra-dy-la-li-a (brad'e-lā'lē-ā). syn bradyarthria. [brady- + G. ialia, speech]

bra-dy-lex-i-a (brad'ē-lek'sē-š). Abnormal slowness in reading. [brady- + G. lexis, word]

bra-dy-10-gi-a (brad'ē-10'jē-ā). syn bradyarthria, [brady- + G. logos, word)

bra-dy-pep-sl-a (brad'ē-pep'sē-š). Slowness of digestion. [brady-+G. pepsis, digestion]

bra-dy-pha-gl-a (brad'ē-fā'jē-ā). Slowness in cating. [brady-+ G. phagō, to cut

bra-dy-pha-si-a (brad'e-fe'ze-ä). A form of aphasia characterized by abnormal slowness of speech. syn bradyphemia. [brady- + G. phasis, speaking]

bra-dy-phe-ml-a (brad'ē-tē'mē-ā). syn brudyphasia. 1brady-+ G. phömē, speech]

bradyphrenia (hrs-de'frā-nā-n). Slowness in mental processing due to a decreased ability to shift quickly from one conceptual pattern to another; most often seen with Parkinson disease. [brady- + -phrenia]

bra-dyp-ne-a (brad-ip-ne'a). In the diphthong pa, the p is silent only at the beginning of a word, Although bradypne'a is the correct pronunciation, the alternative pronunciation bradyp'nea is widespread in the U.S. Abnormal slowness of respitation, specifically a low respiratory frequency. [brady- + G. pnoē, breath-

bra-dy-psy-chi-a (brad'e-si'ke-a). Slowness of mental reactions. [brady- + G. psychē, soul]

bra-dy-rhyth-mi-a (brad'ē-ridh'mē-ā). syn bradycardia. bra-dy-sper-ma-tism (brad'e-sper'ma-tizm). Absence of ejaculatory force, so that the semen trickles away slowly. [brady + G. sperma (spermat-), seed, + ism)

bra-dy-sphyg-mi-a (brad'e-sfig'me-ä). Slowness of the pulse; can occur without bradycardia, as in ventricular bigernin, when every alternate beat may fail to produce a peripheral pulse, [brady- + G. sphygmos, pulse]

bra-dy-stal-sis (brad'e-stahl'sis). Slow bowel motion. [G. bradys, slow, + (peri) stalsis, contracting around]

bra-dy-tel-e-o-kl-ne-si-a (brad'ē-tel'ē-ō-kin-ē'sē-ŭ). Sudden arrest of a movement just before its intended termination, then after a pause it is completed slowly or by jerks; a symptom of cerebellar disease. SYN bradytelcokinesis. [brady-+ G. teleos. complete, + kinēsis, movement]

bra-dy-tel-e-o-ki-ne-sis (brad'ē-tel'ē-ō-ki-në'sis). syn bradyteleokinesia.

bra·dy-u-ri·a (brad'ē-yū'rē-ā). Slow micturition, [brady- + G. *ouron*, ucine)

bra-dy-zo-ite (brad'ē-zō'īc). A slowly multiplying encysted form of sporozoan parasite typical of chronic infection with Toxoplasma gondil. It has also been called a merozoite or zoite; the complex of b.'s within an enclosing membrane has also been called a pseudocyst, though it is now regarded as a true cyst. [brady-+ G. zōē, life]

braille (brail). A system of writing and printing by means of raised dots corresponding to letters, numbers, and punctuation to enable the blind to read by touch. [Louis Bruille, French teacher of blind, 1809-1852)

Brails-ford (brails'ford). James Proderick, English radiologist, 1888-1961, SEE B.-Morquio disease.

Brain (brin), Walter Russell, English physician, 1895-1966, see B. reflex.

Hibrain (brain) [TA]. That part of the central nervous system contained within the cranium, see Also encephalon, Cf. cerebrum. cerebellum. See page 251, B13. [A.S. braegen]

eloquent b., those parts of the b. that control speech, motor functions, and senses, localization of which is important in treating b. tumors.

split b., a b. in which the corpus callosum and usually the anterior and posterior commissures have been sectioned, usually to treat certain refractory epilepsies. visceral b., xvv limbic system.

brain-case (branker). SYN neurocranium.

brain-stem, brain stem (bran'stem) [TA]. Originally, the entire unpaired subdivision of the brain, composed of (in americal sequence) the rhombencephalon, mesencephalon, and diencephalon as distinguished from the brain's only paired subdivision, the telencephalon. More recently, the term's connotation has under gone several arbitrary modifications: some use it to denote a more than rhombencephalon plus mesencephalon, distinguishing that complex from the presencephalon (diencephalon plus teleg-cephalon); others restrict it even further to refer exclusively to the rhombencephalon. From both developmental and architectural viewpoints, the original interpretation seems preferable. syn true cus encephali [TA].

brain-wash-ing (bran'wash'ing). Inducing a person to modify minudes and behavior in certain directions through various forms of psychological pressure or tomure.

bran (bran). A by-product of the milling of wheat, containing approximately 20% of indigestible cellulose; a bulk catharic usually taken in the form of cereal or special bran products.

branch (branch) [TA]. An offshoot: in anatomy, one of the primary divisions of a nerve or blood vessel. A branch. see ramits artery, nerve, vein. syn ramus (1) [TA].

accessory meningeal b., syn prerygomeningeal arrery. accessory meningeal b. of middle meningeal artery. 50 accessory b. of middle meningeal artery.

accessory b. of middle meningeal artery [TA], a backets the middle meningeal or maxillary artery in the infratempt of the middle meningeal or maxillary artery in the infratempt of the middle meningeal or maxillary artery in the infratempt of the middle meningeal or maxillary artery in the infratempt of the middle meningeal artery [TA], a backets are markets as a second of the middle meningeal artery [TA], a backets are markets as a second of the middle meningeal artery [TA], a backets are markets as a second of the middle meningeal artery [TA], a backets are middle meningeal artery in the infratempt of the middle meningeal artery in t ral fossu and passing superiorly through the foreign ovale is supply the trigenimal ganglion, dura mater, and inner table

浅机构外特許事務所

riosteologi teotomy

ln. alvoji lontium et

SYN Pe

M-3117501

condition

ombilie.

onfile /chīưm

· O May

the

x lm Si

apinza)

SUC


ÉÜ

11

inputs the periosteum to the bone. SYN periostotomy.

18 16 OUS (per e-os te-us). syn periosteal.

os ie um, pi. per-l-os-te-a (per ē-os tē-um, -ā) [TA]. fibrous membrane covering the entire surface of a bone listanticular cartilage and the areas where it attaches to and ligaments. In young bones, it consists of two layers:
Allular layer that is osteogenic, forming new bone tissue, if fibrous connective tissue layer conveying the blood and merves supplying the bone; in older bones, the osteo-ing reduced, see also perichondral bone. See this reperiose [Mod. L. fr. G. periosteon, nu. of adj. periostemid bones, fr. peri, around, + osteon, bone]



usolat p., p. alveolare, syn periodontium.

manif [[PA], san pericranium.

eldios ti-tis (per e-os-titis). Inflammation of the perioseum.

Parios-to-ma (per e-os-tō mā). SYN periosteoma.

etilios to sis. pl. per-i-os-to-ses (per'e-os-to'sis, -sez).

OS-105-18-1-115 (per/ē-os/tos-tē-7/tis). Inflammation of a hit-involvement of the periosteum. [periosteum + G. oste-(intermedian)

105-10-10me (per e-os-to-tom), syn periosteotome.

910.11C (per e-o'ik, -or'ik). Surrounding the internal car, the petrous portion of the temporal bone, or the spaces allossees in the bony labyrinth that surround the membranous minip[peri- + G. ous, ear]

(1:0: Va·ri·tis (per ē-ē-vā-rī ds). syn perioophoritis.

O. VU-lar (per z-ō'vyū-lär). Surrounding the ovum.

pach.y.men.in.gi.tis (per i-pak e-men in-ji (is). Inmation of the area between the dura and bony covering of the inilarmation]

Pan:cre-a-ti-tle (pcr'i-pan kre-a-ti'tis). Inflammation Deritoneal coat of the pancreus.

pap·il·lar·y (per'i-pap'i-lar'ë). Surrounding a papilla.

per-i-pa-tet-ic (peri-pa-tetik). 1. Walking around; formerly used to describe a patient with "walking" (i.e., mild) typhoid fever. 2. Relating to a disease imported to a nonendetraic area by a host clinically unaffected during the transport phase. [G. peripatësis, a walking about]

per-i-pe-ni-ai (per'i-pe'ne-il). Surrounding the penis.

per-i-pha-ryn-ge-al (per'i-fa-rin'je-al). Surrounding the pharynx.

pe-riph-er-ad (pe-riffer-ad). In a direction toward the periphery. [G. periphereia, periphery, + L. ad, to]

pe-riph-er-al (pe-rifer-al) [TA]. I. Relating to or situated at the periphery, 2. Situated nearer the periphery of an organ or part of the body in relation to a specific reference point; opposite of central (centralis). syn peripheralis [TA], eccentric (3).

pe-riph-e-ra-lis (pe-rif'e-ra'lis) [TA]. syn peripheral.

per-i-pher-in (perifer-in) [MIM\*170710]. A glycoprotein apparently necessary to maintain the shape of the outer segment disc membranes of rods and cones; it is thought by many investigators that a defect in p. is associated with certain types of blindness

pe-riph-e-ro-cen-tral (pe-rife-ro-sen'tral). Relating to both the periphery and the center of the body or any part.

pe-riph-e-ry (pe-rif's-ri). 1. The part of a body away from the center; the outer part of surface. 2. syn denture border. [G. periphereia, fr. peri, around, + phero, to carty]

per-i-phie-bit-ic (per'i-fle-bit'ik). Relating to periphicbitis.

per-i-phie-bi-tis (per'i-fle-bi tis). Inflammation of the outer coat of a vein or of the tissues surrounding it. [peri- + G. phleps. voin, + -itis, inflammation]

Per I. pla-ne-ta (per i-pla-ne ta). A genus of large cockroaches including several cosmopolitan household posts found wherever food is available, especially in moist protected areas. P. americana (American cockroach), a very large brownish-chestnut species, 30-40 mm long, la probably native to Africa but now universally distributed; P. fuliginosa (the smoky-brown cockroach) is a common household pest in the eastern and southeastern U.S. [peri- 4 G. planētēs, à roamer]

per-i-plasm (peri-plazm). The space between the cell membranes and the cell wall, in gram-negative bacteria; contains proteins secreted by the cell.

pe-rip-lo-cin (pe-rip'lo-sin). A cardiotonic glycoside obtained from the bark and stems of Periploca gracea (family Asclepiadaceae), a plant of southern Europe and Western Asia. [G. peri-plokē, a winding around, fr. plekē, to twine, plait]

per-i-po-lar (per'i-po'liir). Surrounding the pole or poles of any body, or any electric or magnetic poles.

per-i-po-le-sis (per'i-pō-lē'sis). Penetration of migrating cells between fixed dissue cells that are normally in close contact. [peri-+ G. poleomai, to wander]

per-i-por-i-tis (peri-por-i'tis). Miliary papules and papuloves-icles with staphylococcic infection; most frequently on the face and in infants. [peri- + G. poros, pore, + -itis, inflammation]

per-i-por-tal (per'i-pōr'iši). Surrounding the portal vein. SYN peripylic.

per-i-proc-tic (per'i-prok'tik). syn circumanal. [pcri- + G. proktos, anusl

per-i-proc-ti-tis (peri-prok-titis). Inflammation of the arcolar tissue about the rectum. SYN perirectitis.

per-i-pros-tat-ic (per'i-pros-tat'ik). Surrounding the prostate.

per-i-pros-ta-ti-tis (per'i-pros'ti-ti'tis). Obsolete term for inflammation of the tissues surrounding the prostate.

per-i-py-le-phle-bi-tis (peri-prile-fie-bi'tis). Inflammation of the tissues around the portal vein. [peri- + G. pyle, gate, + phleps, vein, + -itis, inflammation]

per-l-py-lic (per'i-pi'lik). Syn periportal. [peri- + G. pyle, portal, gatel

per-i-py-lor-ic (per'i-pī-lōr'ik, -pǐ-lōr'ik). Surrounding the py-